



Atherosclerosis Risk in Communities Study

Cohort Exam Visit 6 NCS

DERIVE62_np Derived Variable Dictionary (v1)

February 2025

ARIC Visit 6 Derived Variable Dictionary

Table of Contents

New or Changed from PREVIOUS Distribution.....	9
1. Overview.....	11
2. Administrative.....	11
2.1 SUBJECTID (ARIC Subject ID (CIR)).....	11
2.2 ID (ARIC ID - same as SUBJECTID).....	11
2.3 CENTER (Field Center).....	11
2.4 V6CENTER (Visit 6 Field Center).....	12
2.5 V6DATE61_FollowUpDays (Days of follow up from visit 1 to Visit 6 Date).....	12
2.6 V6DATE61_year (Year of Visit 6 Date).....	12
2.7 RES_OTH (Restrictions on Other Procedures).....	12
2.8 RES_DNA (Restrictions on DNA Usage).....	13
2.9 LASTFUINTERVIEW_DATE61_FUDays (Days of follow up from visit 1 to Date of last completed follow-up interview).....	13
2.10 LASTFUINTERVIEW_DATE61_year (Year of Date of last completed follow-up interview).....	14
2.11 STAGE_1_TYPE (V6 Type of Stage 1 Exam).....	14
3. Socio-Demographic.....	15
3.1 GENDER (Sex).....	15
3.2 GENDER61 (Corrected Gender (V1CORGE1)).....	15
3.3 RACEGRP (Race).....	15
3.4 RACEGRP61 (Corrected Race (V1CORRA1)).....	15
3.5 BIRTHDAT (Date of Birth).....	16
3.6 BIRTHDAT61 (Corrected Birthdate (V1CORBIR)).....	16
3.7 V6AGE61 (Visit 6 Age).....	17
3.8 V6AGE62 (Corrected Visit 6 Age).....	17

4. Anthropometry and Blood Pressure	18
4.1 BMI61 (V6 Body Mass Index in kg/m ²)	18
4.2 WSTHPR61 (V6 Waist-to-Hip Ratio).....	18
4.3 SYSTOLIC61 (V6 Mean Systolic BP of 2nd and 3rd Measurements)	18
4.4 DIASTOLIC61 (V6 Mean Diastolic BP of 2nd and 3rd Measurements).....	18
4.5 PULSE61 (V6 Mean Pulse of 2nd and 3rd Measurements)	19
5. Alcohol Use.....	20
5.1 DRNKR61 (V6 Drinker Status).....	20
5.2 ETHANL61 (V6 Usual Ethanol Intake (g/wk))	20
5.3 CURDRK61 (V6 Current Drinker)	20
5.4 FORDRK61 (V6 Former Drinker).....	21
5.5 EVRDRK61 (V6 Ever Drinker)	21
6. Smoking.....	22
6.1 CURSMK62 (V6 Current cigarette smoker)	22
6.2 FORSMK62 (V6 Former cigarette smoker).....	22
6.3 EVRSMK62 (V6 Ever smoked cigarettes)	23
6.4 CIGT62 (V6 Cigarette Smoking Status).....	24
7. Disease Prevalence.....	25
7.1 DIABTS63 (V6 Diabetes - Lower Cutpoint 140 mg/dL)	25
7.2 DIABTS64 (V6 Diabetes - Lower Cutpoint 126 mg/dL)	26
7.3 DIABTS65 (V6 Diabetes Lab and Meds Only - Lower Cutpoint 126 mg/dL)	26
7.4 DIABTS66 (V6 Diabetes – Hemoglobin A1C, cutpoint 6.5%).....	27
7.5 DIABTS67 (V6 Diabetes (DM medications or DM reported on AFU))	27
7.6 MDDXMI62 (V6 MD Diagnosed Myocardial Infarction)	28
7.7 HXOFMI62 (V6 History of Myocardial Infarction).....	28
7.8 PRVCHD61 (V6 Prevalent CHD before Visit 6)	30
7.9 PRVCHD63 (V6 Prevalent CHD by end of Visit 6)	30
7.10 PRVCHD64 (V6 Prevalent CHD - unverified)	31
7.11 PREVDEFHF61 (V6 Prevalent Definite Heart Failure for Closed Event Years).....	34

7.12	PREVDEFPOSSH61 (V6 Prevalent Definite OR Possible Heart Failure for Closed Event Years)	35
7.13	PRVSTR61 (V6 Prevalent Stroke by the end of Visit 6)	37
7.14	PRVSTR62 (V6 Prevalent Stroke-unverified)	37
7.15	HYPERT64 (V6 Hypertension, definition 4 (DIASTOLIC61 GE 90 or HTN med))	38
7.16	HYPERT65 (V6 Hypertension, definition 5 (SYSTOLIC61 GE 140 or DIASTOLIC61 GE 90 or HTN medication))	38
7.17	HYPERT66 (V6 Hypertension, definition 6 (SYSTOLIC61 GE 160 or DIASTOLIC61 GE 95 or HTN medication)).....	39
7.18	HYPERT67 (V6 Hypertension, definition 7 (SYSTOLIC61 GE 150 or DIASTOLIC61 GE 90 or HTN medication)).....	39
8.	Medication Use	40
8.1	CHOLMDCODE61 (V6 Cholesterol Lowering Medication in past 4 weeks - Using Medi-Span GPI Code).....	42
8.2	CHOLMDCODE62 (V6 Medications Which Secondarily Affect Cholesterol in past 4 weeks - Using Medi-Span GPI Code)	43
8.3	HYPTMD61 (V6 Hypertension Medications in Past 4 Weeks: Self-reported)	43
8.4	HYPTMDCODE61 (V6 Hypertension Lowering Medication in Past 4 weeks - Using Medi-Span GPI code)	44
8.5	STATINCODE61 (V6 Statin Use in past 4 weeks - Using Medi-Span GPI Code).....	44
8.6	ANTICOAGCODE61 (V6 Anticoagulant Use in past 4 weeks - Using Medi-Span GPI Code)	45
8.7	ASPIRINCODE61 (V6 Aspirin Use in past 4 weeks - Using Medi-Span GPI Code).....	45
8.8	ANTIANXMDCODE61 (V6 Antianxiety Medication in past 4 weeks - Using Medi-Span GPI Code).....	46
8.9	ANTIPSYCHMDCODE61 (V6 Antipsychotic Medication in past 4 weeks - Using Medi-Span GPI Code).....	46
8.10	HYPNOTMDCODE61 (V6 Hypnotic/Sedative Medication in past 4 weeks - Using Medi-Span GPI Code).....	47
8.11	ANTICONVMDCODE61 (V6 Anticonvulsant Medication in past 4 weeks - Using Medi-Span GPI Code).....	47

8.12	ANTIDEMDCODE61 (V6 Antidementia/Nootropic Medication in past 4 weeks - Using Medi-Span GPI Code)	48
8.13	CNSALTMDCODE61 (V6 CNS Altering Medication in past 4 weeks - Using Medi-Span GPI Code)	48
8.14	DIABMDCODE61 (V6 Diabetic Medications in past 4 weeks - Using Medi-Span GPI Code)	49
8.15	BETAMDCODE61 (V6 Beta-Blocker in past 4 weeks - Using Medi-Span GPI Code).....	49
8.16	ANGINHMDCODE61 (V6 Angiotensin converting enzyme inhibitor in past 4 weeks - Using Medi-Span GPI Code)	50
8.17	ANGIANTMDCODE61 (V6 Angiotensin II receptor antagonists in past 4 weeks - Using Medi-Span GPI Code)	50
8.18	ALDANTMDCODE61 (V6 Aldosterone Antagonist in past 4 weeks - Using Medi-Span GPI Code)	51
8.19	LOOPDIUMDCODE61 (V6 Loop Diuretic in past 4 weeks - Using Medi-Span GPI Code)	51
8.20	DIGMDCODE61 (V6 Digoxin in past 4 weeks - Using Medi-Span GPI Code)	51
9.	Physical Activity	53
9.1	SPRT_I61 (V6 Sport during Leisure Time)	53
9.2	LISR_I62 (V6 Physical Activity during Leisure Time Excluding Sport)	55
10.	Physical Function.....	57
10.1	SPPBCS61 (V6 Physical Function Chair Stand)	57
10.2	SPPBST61 (V6 Physical Function Semi Tandem Stand)	57
10.3	SPPBSBS61 (V6 Physical Function Side-by-Side Stand)	57
10.4	SPPBTS61 (V6 Physical Function Tandem Stand)	58
10.5	SPPBBAL61 (V6 Physical Function Summary Balance Score).....	58
10.6	WALK4M61 (V6 Physical Function 4 Meter Walk, Fastest Time of 2 Trials).....	58
10.7	WALKAID61 (V6 Physical Function 4 Meter Walk: Used Walking Aid)	59
10.8	WALK4M62 (V6 Physical Function 4 Meter Walk, Average Time of 2 Trials (BOTH TRIAL WITH AID OR BOTH TRIALS WITHOUT AID))	59
10.9	WALKAID62 (V6 Physical Function 4 Meter Walk: Used Walking Aid in both trials (USE WITH WALK4M62))	59
10.10	SPPB4M61 (V6 Physical Function 4 Meter Walk Score).....	60

10.11	SPPB61 (V6 Short Physical Performance Summary Battery Score).....	60
10.12	GRIPBEST61 (V6 Physical Function Grip, Best of 2 Trials).....	60
10.13	GRIPMEAN61 (V6 Physical Function Grip, Mean of 2 Trials or Result for 1 Trial)	61
10.14	V5V6WTDELTA61 (% of V5 weight change from V5 to V6 (neg val means loss) – frailty weight loss component)	61
10.15	WALKSPEED15FT61 (Time in seconds used to walk 15ft – frailty slowness component)	61
10.16	V6CANCER61 (Reported a cancer site during an ARIC follow-up interview in the time from 06-15-2016 to the V6date61 (V6 visit date)).....	62
10.17	TR1WALK4MSP61 (V6 4M Walking Speed for Trial 1 (m/sec)).....	62
10.18	TR2WALK4MSP61 (V6 4M Walking Speed for Trial 2 (m/sec)).....	62
10.19	AVGWALK4MSP61 (V6 Average 4M Walking Speed (m/sec)).....	63
10.20	MINWALK4MSP61 (V6 Fastest 4M Walking Speed for Both Trials (m/sec)).....	63
10.21	UNABLETOWALK61 (Indicator Variable Noting PPT No Attempt to Walk 4M, Not Able)	63
10.22	UNINTEND_WTLOSS61 (V6 Unintentional Weight Loss)	63
10.23	TMW_TOTFT61 (V6 Total Feet Walked in 2-Minute Walk).....	64
10.24	TMW_COMPSPEEDFTPERSEC61 (V6 TMW Speed for Completers (ft/sec))	64
10.25	TMW_COMPSPEEDMPERSEC61 (V6 TMW Speed for Completers (m/sec)).....	64
10.26	TMW_NONCOMPSECWALK61 (V6 TMW Seconds Walking for Noncompleters).....	64
10.27	TMW_NONCOMPSPEEDFTPERSEC61 (V6 TMW Speed for Noncompleters (ft/sec))	65
10.28	TMW_NONCOMPSPEEDMPERSEC61 (V6 TMW Speed for Noncompleters (m/sec))	65
10.29	PAQ_LIMITEDFUNCTION61 (V6 PAQ Limited Function).....	65
10.30	PAQ_GOODFUNCTION61 (V6 PAQ Good Function).....	66
10.31	EXHAUST61 (V6 Responded 2 or 3 on CES3 or CES11 (CESD) – frailty exhaustion component).....	66
10.32	LOWENERGYCOMP61 (V6 Lowest quintile of SPRT_i61 - frailty low physical activity component)	67

10.33	WTLOSSCOMPA61 (V6 Lost >5% weight or Low BMI (<18.5) – frailty weight loss component for 61a definition)	67
10.34	WTLOSSCOMPB61 (V6 Lost >10% weight or Low BMI (<18.5) – frailty weight loss component for 61b definition)	68
10.35	WALKSPEEDCOMP61 (V6 Slowest 20% time to walk 15ft – frailty slowness component)	68
10.36	GRIPSTRENGTHCOMP61 (V6 Slowest 20% grip strength – frailty weakness component)	69
10.37	FRAILITY61a (V6 ARIC Physical Function WG Frailty Definition (a))	70
10.38	FRAILITY61b (V6 ARIC Physical Function WG Frailty Definition (b))	71
10.39	FRAILITY63 (V6 ARIC Physical Function WG Frailty Definition – weight loss dropped)	72
10.40	FRAILITY64 (V6 ARIC Physical Function WG Frailty Definition (unintentional weight loss))	73
10.41	GAITSPEED61 (V6 Gait Speed (m/sec)).....	74
11.	Hearing Results	75
11.1	PTARIGHT61 (Four Frequency Pure-tone Average (PTA) of the Right Ear (Db HL))	75
11.2	PTALEFT61 (Four Frequency Pure-tone Average (PTA) of the Left Ear (Db HL))	75
11.3	PTABETTER61 (Four Frequency Pure-tone Average (PTA) of the Better Ear (smaller PTA) (Db HL))	75
11.4	PTABETTERCAT61 (Hearing Loss Category Based on the Four Frequency Pure-tone Average (PTA) of the B)	76
12.	Neurocognitive Study.....	77
12.1	CESD61 (V6 CES-Depression Scale)	77
12.2	FAQ61 (Functional Activities Questionnaire)	77
13.	SF-12 OUTCOMES	77
13.1	SF12PF61 (V6 SF-12 Physical Functioning Scale)	78
13.2	SF12RP61 (V6 SF-12 Role Physical Scale)	78
13.3	SF12BP61 (V6 SF-12 Bodily Pain Scale)	78
13.4	SF12GH61 (V6 SF-12 General Health Scale)	78
13.5	SF12VT61 (V6 SF-12 Vitality Scale).....	79

13.6	SF12SF61 (V6 SF-12 Social Functioning Scale)	79
13.7	SF12RE61 (V6 SF-12 Role Emotional Scale)	79
13.8	SF12MH61 (V6 SF-12 Mental Health Scale)	79
13.9	SF12PFZ61 (V6 SF-12 Physical Functioning Scale Z Transformation)	80
13.10	SF12RPZ61 (V6 SF-12 Role Physical Scale Z Transformation)	80
13.11	SF12BPZ61 (V6 SF-12 Bodily Pain Scale Z Transformation)	80
13.12	SF12GHZ61 (V6 SF-12 General Health Scale Z Transformation)	80
13.13	SF12VTZ61 (V6 SF-12 Vitality Scale Z Transformation)	81
13.14	SF12SFZ61 (V6 SF-12 Social Functioning Scale Z Transformation)	81
13.15	SF12REZ61 (V6 SF-12 Role Emotional Scale Z Transformation)	81
13.16	SF12MHZ61 (V6 SF-12 Mental Health Scale Z Transformation)	82
13.17	SF12PFT61 (V6 SF-12 Physical Functioning Scale T Score)	82
13.18	SF12RPT61 (V6 SF-12 Role Physical Scale T Score)	82
13.19	SF12BPT61 (V6 SF-12 Bodily Pain Scale T Score)	83
13.20	SF12GHT61 (V6 SF-12 General Health Scale T Score)	83
13.21	SF12VTT61 (V6 SF-12 Vitality Scale T Score)	83
13.22	SF12SFT61 (V6 SF-12 Social Functioning Scale T Score)	83
13.23	SF12RET61 (V6 SF-12 Role Emotional Scale T Score)	84
13.24	SF12MHT61 (V6 SF-12 Mental Health Scale T Score)	84
13.25	SF12AGGPHYS61 (V6 SF-12 Aggregate Physical Health Score)	84
13.26	SF12AGGMENT61 (V6 SF-12 Aggregate Mental Health Score)	85
13.27	SF12AGGPHYST61 (V6 SF-12 Aggregate Physical Health T Score)	85
13.28	SF12AGGMENTT61 (V6 SF-12 Aggregate Mental Health T Score)	85

NEW OR CHANGED FROM PREVIOUS DISTRIBUTION

This table describes the changes to the last published DERIVE62 dictionary. As the dataset undergoes modifications, this table will describe the updates made to the previously distributed dataset.

Modification Date	Variable Name	Reason(s) for Change
02/14/2024	<p>V6DATE61 LASTFUINTERVIEW_DATE61</p> <p>ALL derived lab variables: FAST0861 FAST1261 GLUSIU61 GLUSIU62 TGLEFH61 TCHSIU61 HDLSIU61 LDLSIU61 TRGSIU61 LDL61 EGFR61 EGFRCR61 EGFRCYSC61</p> <p>Operational neurocognitive variables: SELECTED_STAGE_2 NEUROCOGSTAT61 ALGDX61 REVIEWERSYND61 DERMVTTYPE61 LOWPROMMSE61 LOWMEMDOM61 LOWEFDOM61 LOWLANGDOM61 SIGCOGDECLINE61 BLOCKAONLY GCPBLOCKA61</p> <p>Analytic neurocognitive variables: PRORATEDMMSE62 ALGDXSTRATUM61 COGDIAG61</p>	<p>DERIVE62 has undergone the no PHI transformation which means the date variables have been transformed to days since V1 enrollment.</p> <p>The lab-related derived variables in the dataset have been moved to LONGLABV91_YYMMDD.</p> <p>The operational neurocognitive variables have been moved to a non-distributed dataset.</p> <p>The analytic neurocognitive variables have been moved to STATUS81.</p>
09/27/2019	<p>Added ALL SF12 variables: SF12PF61, SF12RP61, SF12BP61, SF12GH61, SF12VT61, SF12SF61, SF12RE61, SF12MH61, SF12PFZ61, SF12RPZ61, SF12BPZ61, SF12GHZ61, SF12VTZ61, SF12SFZ61, SF12REZ61, SF12MHZ61, SF12PFT61, SF12RPT61, SF12BPT61, SF12GHT61, SF12VTT61, SF12SFT61, SF12RET61, SF12MHT61,</p>	<p>The company that holds the license has instructed us to include the statement "<u>a modified version of the SF-12v2 was used</u>" in ALL manuscripts using SF-12 data.</p> <p>Note that the modifications are very minor, things like underlining certain words or phrases in the questions, and largely</p>

	SF12AGGPHYS61, SF12AGGMENT61, SF12AGGPHYST61, SF12AGGMENTT61	irrelevant because the instrument was interviewer-administered.
09/27/2019	FORSMK62, EVRSMK62, CIGT62, EGFR61, PRVCHD61, PRVCHD63, PRVCHD64, PREVDEFHF61, PREVDEFPOSSH61, PRVSTR61, DEMRVTYPE61, EXHAUST61, FRAILTY61a, FRAILTY61b, FRAILTY63, FRAILTY64, and GAITSPEED61	Added variables that were missing from the previously distributed datasets.
09/27/2019		Derived variables using event year 2017 surveillance data are final and frozen.
9/12/2018	REVIEWERSYND61, COGDIAG61	Updated to reflect the final classification for the 8 adjudicated reviews that were pending at the time of the previous data distribution.
07/18/2018	LASTFUINTERVIEW_DATE61, DIABTS63, DIABTS64, DIABTS66, DIABTS67, MDDXMI62, HXOFMI62, PRVSTR62	Updated the variables that use follow-up data with the most recently distributed composite follow-up dataset (uc750301_compositeafu_safu). See codebook on ARIC website for updated distributions of these variables.
03/16/2018	NEUROCOGSTAT61, ALGDX61, ALGDXSTRATUM61	An error in the specifications for calculating the predicted domain scores for V6 affected the variables NEUROCOGSTAT61, ALGDX61, ALGDXSTRATUM61. This dataset includes corrected values for these variables.

1. OVERVIEW

The DERIVE62 dataset has 4,003 records, one for each participant who completed Stage 1 at Visit 6. The purpose of this dataset is to provide ARIC collaborators widely used, verified derived variables, many of which are consistent with variables derived at prior visits.

The dataset naming conventions are as follows: The dataset name retains the retrieval date (ex: DERIVE61_180718) until the dataset is considered final, frozen. After a dataset is frozen, the retrieval date is dropped from the dataset name (ex: DERIVE61). The first digit in the dataset name refers to the visit number. The second digit in the dataset name is incremented in number when the current dataset undergoes significant changes. The variable naming convention is similar: Across-visit variables have identical names except for the second to last digit in the variable name, which represents the visit number (ex: GENDER51 at Visit 5 vs. GENDER61 at Visit 6). The last digit in the variable name identifies the definition version of a variable.

Most of the variables are derived directly from the data collected at the visit. However, some variables use ARIC cohort surveillance and ARIC follow-up data in their definitions. DERIVE61_yymmdd will be final, frozen after the surveillance datasets are complete for events in 2017. DERIVE62 is final, frozen.

2. ADMINISTRATIVE

2.1 SUBJECTID (ARIC Subject ID (CIR))

Type: Character; length: \$7.

2.2 ID (ARIC ID - same as SUBJECTID)

Description: The historical participant identifier from visits 1-4 is ID. The value of ID is the same value as SUBJECTID. Use ID when merging visit 6/NCS stage 1 data with datasets from previous visits necessary for longitudinal analyses.

Type: Character; length: \$7.

Algorithm: ID=SUBJECTID

Source variable(s): SUBJECTID

2.3 CENTER (Field Center)

Description: Character variable with four possible values derived from the enrollment site:

F: Forsyth County, North Carolina
J: The city of Jackson, Mississippi
M: Selected northwestern suburbs of Minneapolis, Minnesota
W: Washington County, Maryland

Type: Character; length: \$1.

Algorithm: CENTER = First letter of the subject ID

Source variable(s): SUBJECTID

2.4 V6CENTER (Visit 6 Field Center)

Algorithm: The value of V6CENTER is the same as CENTER unless the ARIC study participant has relocated geographically and moved into another field center at visit 6. In that instance, the value of V6CENTER is the value of the field center where the participant was seen.

2.5 V6DATE61_FollowUpDays (Days of follow up from visit 1 to Visit 6 Date)

Description: Variable calculated as the days between V1 enrollment and the V6 clinic visit date. SAS date variable which documents the date of the participant's visit 6 exam. The visit date is determined as the earlier of the date from the Anthropometry form (ANT) or the date from the Sitting Blood Pressure form (SBP). In the instance where 1 of those 2 form dates is missing, the other non-missing date is used.

Type: Numeric

Source variable(s): ANT0a, BIO0a, SBP0a

2.6 V6DATE61_year (Year of Visit 6 Date)

Algorithm: Year determined from MIN(ANT0a, SBP0a)

Type: numeric

Source variable(s): ANT0a, SBP0a

2.7 RES_OTH (Restrictions on Other Procedures)

Description: The derived informed consent file, ICTDER05, includes information and dates of final consents for ARIC participants. Change in consent status is recorded in ARIC follow-up and visit 6 using the ICT form. RES_OTH indicates the types of restriction on other procedures. We

request that the investigators exclude appropriate records with partial restrictions prior to data analysis.

Type: Character; length: \$130.

Algorithm: if ICT3=1 and ICT4=1 and ICT7=1 then res_OTH='Full Consent';
else if ICT3=1 and ICT4=1 and (ICT7=0 or missing(ICT7)) then
res_OTH='Not for Profit';
else if ICT3=1 and ICT4=0 then res_OTH='ARIC Only';
else if ICT3=0 and ICT4=0 then res_OTH='No Consent';

Source variable(s): ICT3, ICT4, ICT7

2.8 RES_DNA (Restrictions on DNA Usage)

Description: The derived informed consent file, ICTDER05, includes information and dates of final consents for ARIC participants. Change in consent status is recorded in ARIC follow-up and visit 6 using the ICT form. The variable RES_DNA indicates the type of restriction on DNA use. We request that the investigators exclude appropriate records with partial restrictions prior to data analysis.

Type: Character; length: \$130.

Algorithm: if ICT5=1 and ICT6=1 and ICT7=1 then RES_DNA='Full Consent';
else if ICT5=1 and ICT6=1 and (ICT7=0 or missing(ICT7)) then
RES_DNA='Not for Profit';
else if ICT5=1 and ICT6=0 then RES_DNA='ARIC Only';
else if ICT5=0 and ICT6=0 then RES_DNA='No use/storage DNA';

Source variable(s): ICT5, ICT6, ICT7

2.9 LASTFUINTERVIEW_DATE61_FUDays (Days of follow up from visit 1 to Date of last completed follow-up interview)

Description: Variable calculated as the days between V1 enrollment and the last AFU follow-up interview where an actual contact was made, prior to January 1, 2018.

Type: Numeric

Algorithm: LASTFUINTERVIEW_DATE61=the max value of AFUcomp1_A in the composite follow-up dataset among the records for a single ID where AFUcomp2_A indicates that the interview was accomplished (AFUcomp2_A in ('A','C','D')) and the date preceded January 1, 2018.

Source variable(s): AFUcomp1_A, AFUcomp2_A

2.10 LASTFUINTERVIEW_DATE61_year (Year of Date of last completed follow-up interview)

Algorithm: Year determined from LASTFUINTERVIEW_DATE61

Type: numeric

Source variable(s): AFUcomp1_A, AFUcomp2_A

2.11 STAGE_1_TYPE (V6 Type of Stage 1 Exam)

Description: Categorical variable that describes the participant's type of Stage 1 exam.

Format: A=Full, B=Abbreviated, C=Home, D=Long Term Care Facility.

Type: Character; length=\$1.

Algorithm: STAGE_1_TYPE=RTS17

Source variable(s): RTS17

3. SOCIO-DEMOGRAPHIC

3.1 GENDER (Sex)

Description: Categorical variable that describes the participant's gender: M=Male, F=Female.

Type: Character; length=\$1.

Algorithm: GENDER = GENDER from DERIVE13

Source variable(s): [DERIVE13] GENDER

3.2 GENDER61 (Corrected Gender (V1CORGE1))

Description: Categorical variable that describes the participant's gender: M=Male, F=Female. Incorrect values for the variable GENDER were identified following the initial data collection on the ARIC cohort. The ARIC Executive Committee has recommended continuing to use the uncorrected variable (GENDER) for Visit 1 and longitudinal analyses. The corrected version could be used for cross-sectional analyses other than Visit 1 and should be decided by the Investigator.

Type: Character; length=\$1.

Algorithm: GENDER61 = V1CORGE1

Source variable(s): V1CORGE1

3.3 RACEGRP (Race)

Description: Categorical variable which describes the participant's race: A=Asian, B=Black, I=Native American, W=White.

Type: Character; length=\$1.

Algorithm: RACEGRP = RACEGRP from DERIVE13

Source variable(s): [DERIVE13] RACEGRP

3.4 RACEGRP61 (Corrected Race (V1CORRA1))

Description: Categorical variable which describes the participant's race: A=Asian, B=Black, I=Native American, W=White. Incorrect values for the variable RACEGRP were identified following the initial data collection on the ARIC cohort. The ARIC Executive Committee has recommended continuing to use the uncorrected variable (RACEGRP) for Visit 1 and longitudinal analyses. The corrected version could be used for cross-sectional analyses other than Visit 1 and should be decided by the Investigator.

Type: Character; length=\$1.

Algorithm: RACEGRP61 = V1CORRA1

Source variable(s): V1CORRA1

3.5 BIRTHDAT (Date of Birth)

Description: SAS date variable that documents the originally reported date of birth for the participant.

Type: Date

Algorithm: BIRTHDAT = BIRTHDAT from DERIVE13

Source variable(s): [DERIVE13] BIRTHDAT

3.6 BIRTHDAT61 (Corrected Birthdate (V1CORBIR))

Description: SAS date variable that corrects known errors in the BIRTHDAT variable. Incorrect values for the variable BIRTHDAT were identified following the initial data collection on the ARIC cohort. The ARIC Executive Committee has recommended continuing to use the uncorrected variable (BIRTHDAT) for Visit 1 and longitudinal analyses. The corrected version could be used for cross-sectional analyses other than Visit 1 and instances where appropriate should be decided by the Investigator.

Type: Date

Algorithm: BIRTHDAT61 = V1CORBIR

Source variable(s): V1CORBIR

3.7 V6AGE61 (Visit 6 Age)

Description: Participant's age at the time of the visit 6 exam calculated from the BIRTHDAT variable.

Type: Numeric

Algorithm: If V6DATE61>.z and BIRTHDAT>.z then V6AGE61 = floor((intck('month', BIRTHDAT,V6DATE61)-(day(V6DATE61) < day(BIRTHDAT)))/12);

Source variable(s): BIRTHDAT (Date of Birth), V6DATE61

3.8 V6AGE62 (Corrected Visit 6 Age)

Description: Participant's age at the time of the visit 6 exam calculated from the BIRTHDAT61 variable. This variable is based on the corrected birthdate. The ARIC Executive Committee has recommended continuing to use the uncorrected variable (V6AGE61) for Visit 1 and longitudinal analyses. The corrected version could be used for cross-sectional analyses other than Visit 1 and should be decided by the Investigator.

Type: Numeric

Algorithm: If V6DATE61 > .z and BIRTHDAT61 > .z then V6AGE62 = floor((intck('month', BIRTHDAT61,V6DATE61) - (day(V6DATE61) < day(BIRTHDAT61)))/12);

Source variable(s): BIRTHDAT61 (Corrected Birthdate), V6DATE61

4. ANTHROPOMETRY AND BLOOD PRESSURE

4.1 BMI61 (V6 Body Mass Index in kg/m²)

Description: Body mass index [Weight (kg)] / [Height (cm) / 100]²

Type: Numeric

Algorithm: If missing (V6IN129) or missing (ANT4) then BMI61=missing;
Else BMI61= ANT4/(V6IN129/100)²

Source variable(s): V6IN129 (Last Measured Height in CM), ANT4

4.2 WSTHPR61 (V6 Waist-to-Hip Ratio)

Description: Ratio of waist girth to hip girth

Type: Numeric

Algorithm: If missing(ANT10a) or missing(ANT10b) or ANT10b=0 then
WSTHPR61=missing
Else WSTHPR61=(ANT10a/ANT10b)

Source variable(s): ANT10a, ANT10b

4.3 SYSTOLIC61 (V6 Mean Systolic BP of 2nd and 3rd Measurements)

Description: Mean of 2nd and 3rd systolic blood pressure measurements, consistent with V1 through V5

Type: Numeric

Algorithm: SYSTOLIC61=mean of SBP8 and SBP11
If SYSTOLIC61=missing then SYSTOLIC61=SBP14

Source variable(s): SBP8, SBP11, SBP14

4.4 DIASTOLIC61 (V6 Mean Diastolic BP of 2nd and 3rd Measurements)

Description: Mean of 2nd and 3rd diastolic blood pressure measurements, consistent with V1 through V5

Type: Numeric

Algorithm: DIASTOLIC61=mean of SBP9 and SBP12
 If DIASTOLIC61=missing then DIASTOLIC61=SBP15

Source variable(s): SBP9, SBP12, SBP15

4.5 PULSE61 (V6 Mean Pulse of 2nd and 3rd Measurements)

Description: Mean of 2nd and 3rd pulse measurements, consistent with V1 through V5

Type: Numeric

Algorithm: PULSE61=mean of SBP10 and SBP13
 If PULSE61=missing then PULSE61=SBP16

Source variable(s): SBP10, SBP13, SBP16

5. ALCOHOL USE

5.1 DRNKR61 (V6 Drinker Status)

Format: 1=Current Drinker
2=Former Drinker
3=Never Drinker
4=Unknown

Type: Numeric

Algorithm: Use the 1st not-permanently missing occurrence of ALC form if more than one.
If ALC2 in ('Y', missing) and ALC3= 'Y' then DRNKR61=1;
Else if (ALC2= 'Y' and ALC3= 'N') then DRNKR61=2;
Else if (ALC2= 'N' and ALC3 in ('N', missing)) then DRNKR61=3;
Else if (ALC2=missing and ALC3= 'N') or (ALC2= 'Y' and ALC3=missing) then DRNKR61=4;
Else DRNKR61=missing.

Source variable(s): ALC2, ALC3

5.2 ETHANL61 (V6 Usual Ethanol Intake (g/wk))

Format: continuous variable

Type: Numeric

Algorithm: Use the 1st not-permanently missing occurrence of ALC form if more than one.
If (DRNKR61 in (2, 3) or ALC3='N') then ETHANL61=0;
Else if (DRNKR61 in (4, missing) and missing(ALC5a) and missing(ALC6a) and missing(ALC7a)) then ETHANL61=missing;
Else ETHANL61=(ALC5a x 10.8) + (ALC6a x 13.2)+(ALC7a x 15.1).

Source variable(s): ALC3, ALC5a, ALC6a, ALC7a, DRNKR61

5.3 CURDRK61 (V6 Current Drinker)

Format: 0=No,
1=Yes,
.T=missing (keeping .T for historical purposes).

Type: Numeric

Algorithm: Use the first not-permanently missing occurrence of ALC form if more than one.
If (ALC2 in ('Y', missing) and ALC3='Y') then CURDRK61=1;
Else if ALC3='N' or (ALC2='N' and ALC3=missing) then CURDRK61=0;
Else CURDRK61=.T

Source variable(s): ALC2, ALC3

5.4 FORDRK61 (V6 Former Drinker)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: Use the first not-permanently missing occurrence of ALC form if more than one.
If (ALC2= 'Y' and ALC3='N') then FORDRK61=1;
Else if (ALC2 in ('Y', missing) and ALC3= 'Y') or (ALC2= 'N' and ALC3 in ('N', missing)) then FORDRK61=0;
Else FORDRK61=.T

Source variable(s): ALC2, ALC3

5.5 EVRDRK61 (V6 Ever Drinker)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: Use the first not-permanently missing occurrence of ALC form if more than one.
If ALC2= 'Y' or (ALC2=missing and ALC3= 'Y') then EVRDRK61=1;
Else if (ALC2= 'N' and ALC3 in ('N', missing)) then EVRDRK61=0;
Else EVRDRK61=.T

Source variable(s): ALC2, ALC3

6. SMOKING

The following definition is a temporary variable used in the algorithms of the smoking derived variables and not found on the DERIVE61 dataset.

Temporary V5V6SMOKECIG (AFU Cigarette Smoker)

Description: derived from composite (AFU) dataset

Algorithm: =1 if any AFUcomp30_G="Y" and AFUcomp1_A is between V5DATE51 (located in DERIVE52) and V6DATE61
=0 if no AFUcomp30_G="Y" and at least one AFUcomp30_G='N' and AFUcomp1_A is between V5DATE51 (located in DERIVE52) and V6DATE61
=missing if all AFUcomp30_G is missing or if V5DATE51 and/or V6DATE61 is missing

The cigarette smoking variables combine responses from the ARIC visit ALC form and ARIC follow-up data.

6.1 CURSMK62 (V6 Current cigarette smoker)

Format: 0=No,
1=Yes,
.T=missing

Type: Numeric

Algorithm: If ALC1= 'Y' then CURSMK62 = 1
Else if ALC1= 'N' then CURSMK62 = 0
Else if ALC1=missing then CURSMK62 = .T

Source variable(s): ALC1

6.2 FORSMK62 (V6 Former cigarette smoker)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If CURSMK62=0 and (CIGT41=1 or CIGT41=2 or CIGT52=1 or CIGT52=2 or V5V6SMOKECIG=1) then FORSMK62=1

Else if CURSMK62=1
OR
(CURSMK62=0 and CIGT41=3 and CIGT52=3 and
V5V6SMOKECIG=0) then FORSMK62=0

Else if (CURSMK62=.T and (CIGT41=1 or CIGT41=2 or CIGT52=1 or
CIGT52=2 or V5V6SMOKECIG=1))
OR
(CURSMK62=0 and CIGT41 is missing and CIGT52 is missing and
V5V6SMOKECIG is missing) then FORSMK62=.T

Else if ALL variables CURSMK62, CIGT41, CIGT52, AND
V5V6SMOKECIG are missing then FORSMK62=.

Source variable(s): CURSMK62, CIGT41, CIGT52, V5V6SMOKECIG

6.3 EVRSMK62 (V6 Ever smoked cigarettes)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If (CURSMK62=1 or CIGT41=1 or CIGT41=2 or CIGT52=1 or
CIGT52=2 or V5V6SMOKECIG=1) then EVRSMK62=1

Else if (CURSMK62=0 and CIGT41=3 and CIGT52=3 and
V5V6SMOKECIG=0) then EVRSMK62=0

Else if (CURSMK62=0 and CIGT41 not 1 and CIGT41 not 2 and
CIGT52 not 1 and CIGT52 not 2 and V5V6SMOKECIG is missing)

OR

(CURSMK62=0 and CIGT41 is missing and CIGT52 is missing and
V5V6SMOKECIG not 1)

OR

(CURSMK62=missing and CIGT41=missing and CIGT52=missing and
V5V6SMOKECIG=0)

OR

(CURSMK62=missing and CIGT41=3 and CIGT52=3 and
V5V6SMOKECIG not 1) then EVRSMK62=.T;

Else if ALL variables CURSMK62, CIGT41, CIGT52, AND
V5V6SMOKECIG are missing then EVRSMK62=.

Source variable(s): CURSMK62, CIGT41, CIGT52, V5V6SMOKECIG

6.4 CIGT62 (V6 Cigarette Smoking Status)

Format: 1=Current smoker,
2=Former smoker,
3=Never smoker,
4=Unknown, but one of the above categories may be ruled out
. =missing.

Type: Numeric

Algorithm: If CURSMK62=1 then CIGT62=1

Else if CURSMK62=0 and (CIGT41=1 or CIGT41=2 or CIGT52=1 or CIGT52=2 or V5V6SMOKECIG=1) then CIGT62=2

Else if CURSMK62=0 and CIGT41=3 and CIGT52=3 and V5V6SMOKECIG=0 then CIGT62=3

Else if (CURSMK62=missing and (CIGT41=1 or CIGT41=2 or CIGT52=1 or CIGT52=2 or V5V6SMOKECIG=1)
OR
(CURSMK62=0 and CIGT41 is missing and CIGT52 is missing and V5V6SMOKECIG is missing) then CIGT62=4

Else if ALL variables CURSMK62, CIGT41, CIGT52, and V5V6SMOKECIG are missing then CIGT62=.

Source variable(s): CURSMK62, CIGT41, CIGT52, V5V6SMOKECIG

7. DISEASE PREVALENCE

There are a number of variables in the Disease Prevalence section that utilize cohort surveillance information, through event year 2017, to derive disease prevalence. Event year 2017 is closed. All derived variables using surveillance data are considered final.

Diabetes Prevalence Variables

Temporary variables were created for use in the algorithms of the following diabetes-derived variables. They are not found on the DERIVE61 dataset.

Temporary MDDX_DIAB61

Algorithm: MDDX_DIAB61=1 if a participant reported being diagnosed during ARIC Follow-Up interviews with diabetes prior to the visit 6 data (V6DATE61). The composite dataset variables considered are AFUCOMP1_A, AFUCOMP7D_G, and AFUCOMP15_M.

Temporary INCSELFREPDM61

Algorithm: If any of the records for a single ID have a 'Y' value for either AFUcomp7d_G or AFUcomp15_M and .z<afucomp1_A<="31DEC2017"d then INCSELFREPDM61 = 1;
Else if AFUcomp7d_G, AFUcomp15_M are (N,") or (" ,N) respectively in all records for a single ID, where .z<afucomp1_A<="31DEC2017"d then INCSELFREPDM61 = 0;
Else INCSELFREPDM61 = .

7.1 DIABTS63 (V6 Diabetes - Lower Cutpoint 140 mg/dL)

Description: Diabetes variable defined as present if glucose value \geq 140 or non-fasting glucose value \geq 200 or using medication for diabetes or self-report diagnosis of diabetes.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If (BIO7a = 'Y' and CHEM5 \geq 140) or (BIO7a ^= 'Y' and CHEM5 \geq 200) or (MSRF2 ^= 'T' and MSRF33c = 'Y') or MDDX_DIAB=1 then DIABTS63=1
Else if (.z<CHEM5<140) and MDDX_DIAB ^= 1 and MSRF33c ^= 'Y' then DIABTS63=0

Else DIABTS63=.T

Source variable(s): BIO7a, CHEM5, MDDX_DIAB, MSRF2, MSRF33c

7.2 DIABTS64 (V6 Diabetes - Lower Cutpoint 126 mg/dL)

Description: Diabetes variable defined as present if glucose value \geq 126 or non-fasting glucose value \geq 200 or using medication for diabetes or self-report diagnosis of diabetes.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If (BIO7a = 'Y' and CHEM5 \geq 126) or (BIO7a ^= 'Y' and CHEM5 \geq 200) or (MSRF2 ^= 'T' and MSRF33c = 'Y') or MDDX_DIAB=1 then DIABTS64=1
Else if (.z<CHEM5<126) and MDDX_DIAB ^= 1 and MSRF33c ^= 'Y' then DIABTS64=0
Else DIABTS64=.T

Source variable(s): BIO7a, CHEM5, MDDX_DIAB, MSRF2, MSRF33c

7.3 DIABTS65 (V6 Diabetes Lab and Meds Only - Lower Cutpoint 126 mg/dL)

Description: Diabetes variable defined as present if glucose value \geq 126 or non-fasting glucose value \geq 200 or using medication for diabetes.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If (BIO7a = 'Y' and CHEM5 \geq 126) or (BIO7a ^= 'Y' and CHEM5 \geq 200) or (MSRF2 ^= 'T' and MSRF33c = 'Y') then DIABTS65=1
Else if (.z<CHEM5<126) and MSRF33c ^= 'Y' then DIABTS65=0
Else DIABTS65=.T

Source variable(s): BIO7a, CHEM5, MSRF2, MSRF33c

7.4 DIABTS66 (V6 Diabetes – Hemoglobin A1C, cutpoint 6.5%)

Description: Diabetes variable defined as present if hemoglobin A1C value $\geq 6.5\%$ or using medication for diabetes or self-report diagnosis of diabetes.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If (CHEM1 ≥ 6.5) or (MSRF2 \neq 'T' and MSRF33c = 'Y') or MDDX_DIAB=1 then DIABTS66=1
Else if (.z < CHEM1 < 6.5) and MDDX_DIAB \neq 1 and MSRF33c \neq 'Y' then DIABTS66=0;
Else DIABTS66=.T.

Source variable(s): CHEM1, MDDX_DIAB, MSRF2, MSRF33c

7.5 DIABTS67 (V6 Diabetes (DM medications or DM reported on AFU))

Description: Diabetes variable defined as present if participant is using medication for diabetes or self-report diagnosis of diabetes.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If (MSRF2 \neq 'T' and MSRF33c='Y') or INCSELFREPDM61=1 or DIABMDCODE61=1 then DIABTS67=1
Else if INCSELFREPDM61 \neq 1 and MSRF33c \neq 'Y' and DIABMDCODE61 \neq 1 then DIABTS67=0
Else DIABTS67=.T

Source variable(s): MSRF2, MSRF33c, [STATUS61] INCSELFREPDM61, DIABMDCODE61

Myocardial Infarction (MI) Prevalence Variables

7.6 MDDXMI62 (V6 MD Diagnosed Myocardial Infarction)

Description: This variable is derived from ARIC Follow-Up questions that ask if the participant was told by a doctor whether or not they'd had a heart attack between visit 5 and visit 6.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If (AFUCOMP7a_G = 'Y' OR AFUCOMP11a_M = 'Y') AND V5DATE51 < AFUCOMP1_A (AFU DATE) < V6DATE61 then MDDXMI62=1
Else if all values of (AFUCOMP7a_G, AFUCOMP11a_M) for records between visit 5 date and visit 6 date are one of the following combinations (",U), (","), (U,") then MDDXMI62=.T
Else MDDXMI62=0

Source variable(s): AFUCOMP7a_G, AFUCOMP11a_M, V5DATE51, V6DATE61, AFUCOMP1_A

7.7 HXOFMI62 (V6 History of Myocardial Infarction)

Description: This variable is derived from ARIC Follow-Up questions that ask if the participant was told by a doctor whether or not they'd had a heart attack as well as questions asking the participant if they'd been hospitalized for heart attack. The follow-up records from the ARIC Follow-Up composite dataset considered for this variable were collected before the end of V6 data collection (31DEC2017).

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If MDDXMI62=1 or (AFUcomp7_V1= 'Y') or (AFUcomp30_deceased_A='Y') or (AFUcomp30non_deceased_A='Y') then HXOFMI62=1
Else if over all records for a single ID the following value combinations are found for (MDDXMI62, AFUcomp7_V1,

AFUcomp30_deceased_A, AFUcomp30non_deceased_A): (.T, "", "", ""),
(.T, "", 'U', ""), (.T, "", "", 'U') then HXOFMI62=.T
Else HXOFMI62=0

Source variable(s): MDDXMI62, AFUCOMP7_V1, AFUCOMP30_deceased_A,
AFUcomp30non_deceased_A

Coronary Heart Disease (CHD) Prevalence Variables

7.8 PRVCHD61 (V6 Prevalent CHD before Visit 6)

Description: This variable is derived from the baseline status of CHD (PRVCHD05) and the closed event years of ARIC Cohort Surveillance data through 2017, where the events occurred prior to the participant's Visit 6.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: PRVCHD61 = 1 if PRVCHD05 = 1 or (C7_IN_17SP = 1 and
. < C7_DATEISP < V6DATE61) or (C7_IN_17SP = 1 and
V6DATE61 = . and C7_DATEISP <= '15JUN2016'd)

PRVCHD61 = 0 if PRVCHD05 = 0 and ((C7_IN_17SP = 0 or
C7_DATEISP >= V6DATE61 > .) or (V6DATE61 = . and
C7_DATEISP > '15JUN2016'd))

ELSE PRVCHD61 = .T

Source variable(s): PRVCHD05, C7_IN_17SP, C7_DATEISP, V6DATE61

7.9 PRVCHD63 (V6 Prevalent CHD by end of Visit 6)

Description: This variable is derived from the baseline status of CHD (PRVCHD05) and the closed event years of ARIC Cohort Surveillance data through 2017, where the events occurred prior to 31DEC2017.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: PRVCHD63 = 1 if PRVCHD05 = 1 or (C7_IN_17SP = 1 and
. < C7_DATEISP < '31DEC2017'd)

PRVCHD63 = 0 if PRVCHD05 = 0 and ((C7_IN_17SP = 0 or
C7_DATEISP >= '31DEC2017'd)

ELSE PRVCHD63 = .T

Source variable(s): PRVCHD05, C7_IN_17SP, C7_DATEISP

7.10 PRVCHD64 (V6 Prevalent CHD - unverified)

Description: This variable is derived from self-reported ARIC Follow-Up data including questions on doctor told participant about heart attack, coronary bypass, and coronary angioplasty on records collected up through 31DEC2017.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: use AFUcomp: only for those records where .z < AFUcomp1_A <= '31DEC2017'd

PRVCHD64 = 1 if MDDXMI62 = 1 or (AFUcomp13a_G = 'Y') or (AFUcomp15a_G = 'Y')

PRVCHD64 = . if over all records for a single ID the following value combinations are found for (MDDXMI62, AFUcomp13a_G, AFUcomp15a_G):(.M or .), (" or 'U'), (" or 'U')

ELSE PRVCHD64 = 0

Source variable(s): AFUcomp1_A, MDDXMI62, AFUcomp13a_G, AFUcomp15a_G

Heart Failure (HF) Prevalence Variables

The heart failure (HF) variables included in this derived variable dictionary define HF prevalence at Visit 6. When conducting analyses of incident HF, it is important to note that the incident variable is available in the ARIC Cohort Surveillance dataset. The variable C7_INCHFxx uses V1 as the baseline and considers an event as a hospitalization with an ICD-9 discharge diagnosis code of 428.x in any position.

The following definitions are temporary variables for use in the algorithms of the heart failure prevalence derived variables and are not included in DERIVE61.

Temporary HospHF_2005onward

Algorithm: For HFC17OCC1 records where
01JAN2005<=HFEVTDATE<V6DATE61
HospHF_2005onward=1, if CHFDIAG in ('A' 'B' 'C') for any record for SubjectID
HospHF_2005onward=0, otherwise

Temporary OnsetDate

Algorithm: Using PHF_180425 and PHFA1104,
If PHF2c is not missing, OnsetDate=PHF2c
Else if PHF7 is not missing, OnsetDate=PHF7
Else if PHFA2c is not missing, OnsetDate=PHFA2c
Else if PHFA7 is not missing, OnsetDate=PHFA7
Else if PHF0a is not missing, OnsetDate=PHF0a
Else if ENTRY_DA is not missing, OnsetDate=ENRTY_DA

Temporary HF_byPHF

Algorithm: Where missing<OnsetDate<V6DATE61
HF_byPHF=1, if phf1='Y' or phfa1='Y'
HF_byPHF=0, otherwise
Keep last record for each SubjectID only after sorted by SubjectID, HF_byPHF, and OnsetDate

Temporary HF_first428_pre2005

Algorithm: Using C17CELB1 records where missing<CELB04<01JAN2005
HF_first428_pre2005=1, if ICD code in first position (CELB10a) is 428.x for any record per SubjectID
HF_first428_pre2005=0, otherwise

Temporary HF_non1st428_pre2005

Algorithm: Using C17CELB1 records where missing<CELB04<01JAN2005
HF_non1st428_pre2005=1, if ICD code is 428.x anywhere but the first position (CELB10b-CELB10z) for any record for the SubjectID
HF_non1st428_pre2005=0, otherwise

Temporary srHF/srHFdt/Form

Algorithm: In a temporary dataset

For [v3]phxa04, srHF=phxa8j, srHFdt=phxa63, Form=NULL

For [v4]phxb04, srHF=phxb6b, srHFdt=phxb21, Form=NULL

For afu&y1104new (y=g, h, i, j, k), srHF=afu&y7b, srHFdt=afu&y1, Form=NULL

For afu&y1104new (y=l, m), srHF='Y' if afu&y7, afu&y8, afu&y9, or afu&y10 is 'Y'; else srHF='N' if afu&y7, afu&y8, afu&y9, or afu&y10 is 'N'.

srHFdt=afu&y1 if not missing; else srHFdt=ENTRY_DA

Form=NULL

For afu_180425, srHF='Y' if afu30 or afu31 is 'Y'; else srHF='N' if afu30 or afu31 is 'N'.

srHFdt=AFU0a if not missing; else srHFdt=DATESTAMPINITIAL_AFU

Form=NULL

For saf_180425, srHF='Y' if saf4, saf4a, saf5, or saf5a is 'Y'; else srHF='N' if saf4, saf4a, saf5, or saf5a is 'N'

srHFdt=saf0a if not missing; else srHFdt=DATESTAMPINITIAL_SAF

Form=NULL

For phfa1104, srHF=phfa1, Form=form

srHFdt is first completed date from the list phfa7, phfa9, ENTRY_DA

For phf_180425, srHF=phf1, Form=form

srHFdt is first completed date from the list phf7, phf9, DATESTAMPINITIAL_PHF

Keep only records where srHF is 'Y' or 'N'

Temporary selfreportHF

Algorithm: From previous temporary dataset, sort by SubjectID, srHFDT, srHF
Keep records where missing<srHFdt<=v6date61 OR Form='PHF'

Start with selfreport=0 for first record per SubjectID then

Selfreport=1, if srHF='Y';

Selfreport=0, if srHF='Y' is followed by a record with srHF='N' and form='PHF'

Last record is kept for each SubjectID

Temporary srHFmed/srHFmedDt

%macro selfreportHFmeds(dsn, med,medDt)

If &med='Y' then do;

srHFmed=1;

srHFmedDt=&medDt;

%mend;

%selfreportHFmeds(v1.msra, MSRA11, msra08d);

%selfreportHFmeds(v2.msrb, MSRB27, msrb24d);

%selfreportHFmeds(v3.msrd04, MSRC29, msrd24e);
 %selfreportHFmeds(v4.msrd04, MSRD31, msrd24e);
 %selfreportHFmeds(v5.msr, MSRF0A, msrf33h);
 %selfreportHFmeds(v6.msr, MSRF0A, msrf33h);
 %selfreportHFmeds(postv4.aful1104new, AFUL1, AFUL46D);
 %selfreportHFmeds(postv4.afum1104new, AFUM1, AFUM46D);
 %selfreportHFmeds(afu.afu_180425, AFU0a, AFU65D);

Temporary selfreportHFmeds

Algorithm: Keep records where missing<srHFmedDT<V6DATE61
 selfreportHFmeds=1, if srHFmed=1 for any record;
 selfreportHFmeds=0, otherwise

7.11 PREVDEFHF61 (V6 Prevalent Definite Heart Failure for Closed Event Years)

Description: A participant is defined to have prevalent definite heart failure if at least one of the following is true: (1) Prior hospitalization (01/01/2005 onward but before V6 visit) classified as Definite (A), Probable (B), or Chronic (C) HF; OR (2) Physician Heart Failure (PHF) Survey with HF onset date prior to V6 (from those with self-reported HF) in which the physician answers YES to "has this patient ever had HF or CM?"; OR (3) Hospitalization with an ICD code 428.x in first position (before 01/01/2005)

Format: 0=No,
 1=Yes,
 .=missing.

Algorithm:

1. Prior hospitalization (01/01/2005 onward but before V6 visit) classified as Definite (A), Probable (B), or Chronic (C) HF
2. Physician Heart Failure (PHF) Survey with HF onset date prior to V6 (from those with self-reported HF) in which the physician answers YES to "has this patient ever had HF or CM?"
3. Hospitalization with an ICD code 428.x in first position (before 01/01/2005)

Using Temporary Variables:
 If V6DATE61 is not missing AND (HospHF_2005onward=1 OR HF_byPHF=1 OR HF_first428_pre2005=1) then PREVDEFHF61=1
 Else PREVDEFHF61=0

Source variables: V6DATE61, HospHF_2005onward, HF_byPHF, HF_first428_pre2005

7.12 PREVDEFPOSSH61 (V6 Prevalent Definite OR Possible Heart Failure for Closed Event Years)

Description: A participant is defined to have prevalent definite OR possible heart failure if at least one of the following is true: (1) Prior hospitalization (01/01/2005 onward but before V6 visit) classified as Definite (A), Probable (B), or Chronic (C) HF; OR (2) Physician Heart Failure Survey with HF onset date prior to V6 (from those with self-reported HF) in which the physician answers YES to "has this patient ever had HF or CM?"; OR (3) Hospitalization with an ICD code 428.x in first position (before 01/01/2005); OR (4) Hospitalization with an ICD code 428.x in any position other than the first position (any time before 01/01/2005); OR (5) Self-report of HF at AFU prior to V6 or at visits 3-4*, not refuted by the physicians health survey (temporal association will need to be considered); OR (6) Self-report of treatment for HF from any study visit or AFU prior to V6. *Note that self-reported HF was only asked at V3 and V4.

Format: 0=No,
1=Yes,
.=missing.

Algorithm:

1. Prior hospitalization (01/01/2005 onward but before V6 visit) classified as Definite (A), Probable (B), or Chronic (C) HF
2. Physician Heart Failure Survey with HF onset date prior to V6 (from those with self-reported HF) in which the physician answers YES to "has this patient ever had HF or CM?"
3. Hospitalization with an ICD code 428.x in first position (before 01/01/2005)
4. Hospitalization with an ICD code 428.x in any position other than the first position (any time before 01/01/2005)
5. Self-report of HF at AFU prior to V6 or at visits 3-4*, not refuted by the physicians health survey (temporal association will need to be considered)
6. Self-report of treatment for HF from any study visit or AFU prior to V6

*Self-reported HF at V1, V2, V5, and V6 not asked

Using Temporary Variables:
If V6DATE61 is not missing AND (PREVDEFHF61=1 OR
HF_non1st428_pre2005=1 OR selfreportHF=1 OR
selfreportHFmeds=1) then PREVDEFPOSSH61=1
Else PREVDEFPOSSH61=0

Source variables: V6DATE61, PREVDEFHF61, HF_non1st428_pre2005, selfreportHF,
selfreportHFmeds

Stroke Prevalence Variables

7.13 PRVSTR61 (V6 Prevalent Stroke by the end of Visit 6)

Description: This variable is derived from the baseline status of stroke (HOM10D) and the closed event years of ARIC Surveillance data on the cohort.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: PRVSTR61 = 1 if HOM10D = "Y" or (C7_IN17DP = 1 and . < C7_ED17DP <= V6DATE61) or (C7_IN17DP = 1 AND V6DATE61 = . and C7_ED17DP <= '31DEC2017'd)

PRVSTR61 = 0 if HOM10D = "N" and ((C7_IN17DP = 0 or C7_ED17DP > V6DATE61 > .) or (V6DATE61 = . and C7_ED17DP > '31DEC2017'd))

Else PRVSTR61 = .T

Source variable(s): HOM10D, C7_IN17DP, C7_ED17DP, V6DATE61

7.14 PRVSTR62 (V6 Prevalent Stroke-unverified)

Description: This variable is derived from self-reported ARIC Follow-Up data including questions that doctor told participant about stroke or TIA or participant hospitalized for stroke on records collected up through 31DEC2017.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If any record within a single ID has a "Y" value is found either AFUcomp29_A or AFUcomp8b_K then PRVSTR62=1
Else if over all records within a single ID the following value combinations are found for (AFUcomp29_A, AFUcomp8b_K): (","), ("U), (U,") then PRVSTR62=missing
Else PRVSTR62=0

Source variable(s): AFUcomp29_A, AFUcomp8b_K

Hypertension Prevalence Variables

7.15 HYPERT64 (V6 Hypertension, definition 4 (DIASTOLIC61 GE 90 or HTN med))

Description: Hypertension definition defined as diastolic blood pressure (mean of 2nd and 3rd measures) ≥ 90 or medication is being taken for high blood pressure.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If (DIASTOLIC61 ≥ 90) or (MSRF2 ^= 'T' and MSRF33d = 'Y') then HYPERT64 = 1
Else if (0 < DIASTOLIC61 < 90) and (MSRF33d = 'N' or (MSRF33d = missing and MSRF2 = 'T')) then HYPERT64=0
Else HYPERT64 = missing

Source variable(s): DIASTOLIC61, MSRF2, MSRF33d

7.16 HYPERT65 (V6 Hypertension, definition 5 (SYSTOLIC61 GE 140 or DIASTOLIC61 GE 90 or HTN medication))

Description: Hypertension is defined as systolic blood pressure (mean of 2nd and 3rd measures) ≥ 140 or diastolic blood pressure (mean of 2nd and 3rd measures) ≥ 90 or medication is being taken for high blood pressure.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If (DIASTOLIC61 ≥ 90) or (SYSTOLIC61 ≥ 140) or (MSRF2 ^= 'T' and MSRF33d = 'Y') then HYPERT65 = 1
Else if (0 < DIASTOLIC61 < 90) and (0 < SYSTOLIC61 < 140) and (MSRF33d = 'N' or (MSRF33d = missing and MSRF2 = 'T')) then HYPERT65 = 0
Else HYPERT65 = .T

Source variable(s): DIASTOLIC61, SYSTOLIC61, MSRF2, MSRF33d

7.17 HYPERT66 (V6 Hypertension, definition 6 (SYSTOLIC61 GE 160 or DIASTOLIC61 GE 95 or HTN medication))

Description: Hypertension is defined as systolic blood pressure (mean of 2nd and 3rd measures) ≥ 160 or diastolic blood pressure (mean of 2nd and 3rd measures) ≥ 95 or medication is being taken for high blood pressure.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If (DIASTOLIC61 ≥ 95) or (SYSTOLIC61 ≥ 160) or (MSRF2 \neq 'T' and MSRF33d = 'Y') then HYPERT66 = 1
Else if (0 < DIASTOLIC61 < 95) and (0 < SYSTOLIC61 < 160) and (MSRF33d = 'N' or (MSRF33d = missing and MSRF2 = 'T')) then HYPERT66 = 0
Else HYPERT66 = .T

Source variable(s): DIASTOLIC61, SYSTOLIC61, MSRF2, MSRF33d

7.18 HYPERT67 (V6 Hypertension, definition 7 (SYSTOLIC61 GE 150 or DIASTOLIC61 GE 90 or HTN medication))

Description: Hypertension is defined as systolic blood pressure (mean of 2nd and 3rd measures) ≥ 150 or diastolic blood pressure (mean of 2nd and 3rd measures) ≥ 90 or medication is being taken for high blood pressure.

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If (DIASTOLIC61 ≥ 90) or (SYSTOLIC61 ≥ 150) or (MSRF2 \neq 'T' and MSRF33d = 'Y') then HYPERT67 = 1
Else if (0 < DIASTOLIC61 < 90) and (0 < SYSTOLIC61 < 150) and (MSRF33d = 'N' or (MSRF33d = missing and MSRF2 = 'T')) then HYPERT67 = 0
Else HYPERT67 = .T

Source variable(s): DIASTOLIC61, SYSTOLIC61, MSRF2, MSRF33d

8. MEDICATION USE

The following definitions are temporary variables, defined here as reference for the following medication-related, derived variables. They are not found on the DERIVE61 dataset.

Temporary ALLMISS

Algorithm: Do over MSRF#a2 where # = 5 to 29:
If any MSRF#a2 >" then ALLMISS=0, else ALLMISS=1

Temporary MSRF#a2_sub10:

Algorithm: Do over MSRF#a2 where # = 5 to 29:
MSRF#a2_sub10=substr(MSRF#a2,1,10)

Temporary MSRF#a2_sub6:

Algorithm: Do over MSRF#a2 where # = 5 to 29:
MSRF#a2_sub6=substr(MSRF#a2,1,6)

Temporary MSRF#a2_sub4:

Algorithm: Do over MSRF#a2 where # = 5 to 29:
MSRF#a2_sub4=substr(MSRF#a2,1,4)

Temporary MSRF#a2_sub2:

Algorithm: Do over MSRF#a2 where # = 5 to 29:
MSRF#a2_sub2=substr(MSRF#a2,1,2)

Temporary FOUNDCHOL61:

Algorithm: Do over MSRF#a2_sub6 where # = 5 to 29:
If MSRF#a2_sub6 in (771030,390000-399999) then
FOUNDCHOL61=1, else FOUNDCHOL61=0.

Temporary FOUNDCHOL62:

Algorithm: Do over MSRF#a2_sub6 where # = 5 to 29:
If MSRF#a2_sub6 in (331000, 332000, 340000, 363000, 369920,
372000, 376000, 379900, 379910) then FOUNDCHOL62=1, else
FOUNDCHOL62=0.

Temporary FOUNDHYPT61:

Algorithm: Do over MSRF#a2_sub6 where # = 5 to 29:
If MSRF#a2_sub6 in (330000-339999 or 340000-349999 or 360000-
369999 or 370000-379999) or (MSRF33d = 'Y') then
FOUNDHYPT61=1, else FOUNDHYPT61=0.

Temporary FOUNDSTAT61:

Algorithm: Do over MSRF#a2_sub4 where # = 5 to 29:
If MSRF#a2_sub4 equal 3940 then FOUNDSTAT61=1, else
FOUNDSTAT61=0.

Temporary FOUNDACOAG61:

Algorithm: Do over MSRF#a2_sub2 where # = 5 to 29:
If MSRF#a2_sub2 equal 83 then FOUNDACOAG61=1, else
FOUNDACOAG61=0.

Temporary FOUNDASP61:

Algorithm: Do over MSRF#a2_sub4 where # = 5 to 29:
If MSRF#a2_sub4 IN (6410, 6499, 6599, 6420) then
FOUNDASP61=1, else FOUNDASP61=0.

Temporary FOUNDANTIANX61:

Algorithm: Do over MSRF#a2_sub6 where # = 5 to 29:
If MSRF#a2_sub6 in (570000, 571000, 571020, 572000) then
FOUNDANTIANX61=1, else FOUNDANTIANX61=0.

Temporary FOUNDANTIPSYCH61:

Algorithm: Do over MSRF#a2_sub6 where # = 5 to 29:
If MSRF#a2_sub6 in (590000, 590700, 591000, 591500, 592000,
593000, 594000, 595000) then FOUNDANTIPSYCH61=1,
FOUNDANTIPSYCH61=0.

Temporary FOUNDHYPNOT61:

Algorithm: Do over MSRF#a2_sub6 where # = 5 to 29:
If MSRF#a2_sub6 in (600000, 601000, 602000, 602010, 602040,
603000, 603099, 609900) then FOUNDHYPNOT61=1,
FOUNDHYPNOT61=0.

Temporary FOUNDANTICONV61:

Algorithm: Do over MSRF#a2_sub6 where # = 5 to 29:
If MSRF#a2_sub6 in (720000, 721000, 721200) then
FOUNDANTICONV61=1, FOUNDANTICONV61=0.

Temporary FOUNDANTIDEM61:

Algorithm: Do over MSRF#a2_sub6 where # = 5 to 29:
If MSRF#a2_sub6 in (620000, 620500, 620510, 620540) then
FOUNDANTIDEM61=1, FOUNDANTIDEMT61=0.

Temporary FOUNDCNSALT61:

Algorithm: Do over MSRF#a2_sub6 where # = 5 to 29:
If MSRF#a2_sub6 in (571000, 932000, 590000, 590700, 591000,
591500, 592000, 593000, 594000, 595000, 601000, 602000, 602010)
then FOUNDCNSALT 61=1, FOUNDCNSALT 61=0.

Temporary FOUNDDIAB61:

Algorithm: Do over MSRF#a2_sub2 where # = 5 to 29:
If MSRF#a2_sub2=27 then FOUNDDIAB61=1, FOUNDDIAB61=0.

Temporary FOUNDBETA61:

Algorithm: Do over MSRF#a2_sub2 and MSRF#a2 where # = 5 to 29:
If MSRF_a2_sub2{i}=33 AND MSRF_a2_sub10{i}^=3310005010 then
FOUNDBETA61=1, else FOUNDBETA61=0.

Temporary FOUNDANGINH61:

Algorithm: Do over MSRF#a2_sub6 where # = 5 to 29:
If MSRF_a2_sub6{i}=361000 then FOUNDANGINH61=1; else
FOUNDANGINH61=0.

Temporary FOUNDANGIANT61:

Algorithm: Do over MSRF#a2_sub6 where # = 5 to 29:
If MSRF_a2_sub6{i}=361500 then FOUNDANGIANT61=1; else
FOUNDANGIANT61=0.

Temporary FOUNDALDANT61:

Algorithm: Do over MSRF#a2 where # = 5 to 29:
If MSRF_a2_sub10{i} IN (3625000000, 3750002000, 3799000220)
then FOUNDALDANT61=1; else FOUNDALDANT61=0.

Temporary FOUNDLOOPDIU61:

Algorithm: Do over MSRF#a2_sub6 where # = 5 to 29:
If MSRF_a2_sub6{i}=372000 then FOUNDLOOPDIU61=1; else
FOUNDLOOPDIU61=0.

Temporary FOUNDDIG61:

Algorithm: Do over MSRF#a2 where # = 5 to 29:
If MSRF_a2{i}=3120001000 then FOUNDDIG61=1; else
FOUNDDIG61=0.

8.1 CHOLMDCODE61 (V6 Cholesterol Lowering Medication in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: (From UC4735 – V4)
If FOUNDCHOL61=1 then CHOLMDCODE61=1

Else if FOUNDCHOL61=0 AND ((ALLMISS=1 AND MSRF2= 'T') OR (ALLMISS=0)) then CHOLMDCODE61=0

Else if ALLMISS=1 AND (MSRF2= 'F' OR MSRF2='') then CHOLMDCODE61=.T

Else CHOLMDCODE61=.

Source variable(s): FOUNDCHOL61, ALLMISS, MSRF2

8.2 CHOLMDCODE62 (V6 Medications Which Secondarily Affect Cholesterol in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: (From UC4735 – V4)
If FOUNDCHOL62=1 then CHOLMDCODE62=1

Else if FOUNDCHOL62=0 AND ((ALLMISS=1 AND MSRF2= 'T') OR (ALLMISS=0)) then CHOLMDCODE62=0

Else if ALLMISS=1 AND (MSRF2= 'F' OR MSRF2='') then CHOLMDCODE62=.T

Else CHOLMDCODE62=.

Source variable(s): FOUNDCHOL62, ALLMISS, MSRF2

8.3 HYPTMD61 (V6 Hypertension Medications in Past 4 Weeks: Self-reported)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If (MSRF2 NE 'T') and (MSRF33D='Y') then HYPTMD61=1;
Else if (MSRF2='T' and MSRF33D='') OR MSRF33D='N' then HYPTMD61 = 0;

Else If ((MSRF2 NE 'T') and (MSRF33D='U' or MSRF33D = '')) or
((MSRF2='T') and (MSRF33D = 'Y' or MSRF33D='U')) then
HYPTMD61=.T;

Source variable(s): MSRF2, MSRF33D

8.4 HYPTMDCODE61 (V6 Hypertension Lowering Medication in Past 4 weeks - Using Medi-Span GPI code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: (From UC4688 – V4)
If FOUNDHYP61=1 OR (MSRF2 NE 'T') and (MSRF33D='Y')) then
HYPTMDCODE61=1

Else if FOUNDHYP61=0 AND ((ALLMISS=1 AND MSRF2= T') OR
(ALLMISS=0) OR (MSRF2= 'T' and MSRF33D='') OR (MSRF33D=
'N')) then HYPTMDCODE61=0

Else if ALLMISS=1 AND (MSRF2= 'F' OR MSRF2='') then
HYPTMDCODE61=.T

Else HYPTMDCODE61=.

Source variable(s): FOUNDHYP61, ALLMISS, MSRF2, MSRF33D

8.5 STATINCODE61 (V6 Statin Use in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: (From UC4892 – V4)
If FOUNDSTAT61=1 then STATINCODE61=1

Else if FOUNDSTAT61=0 AND ((ALLMISS=1 AND MSRF2='T') OR
(ALLMISS=0)) then STATINCODE61=0

Else if ALLMISS=1 AND (MSRF2='F' OR MSRF2='') then
STATINCODE61=.T

Else STATINCODE61=.

Source variable(s): FOUNDSTAT61, ALLMISS, MSRF2

8.6 ANTICOAGCODE61 (V6 Anticoagulant Use in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: (From UC4892 – V4)
If FOUNDACOAG61=1 then ANTICOAGCODE61=1

Else if FOUNDACOAG61=0 AND ((ALLMISS=1 AND MSRF2= 'T')
OR (ALLMISS=0)) then ANTICOAGCODE61=0

Else if ALLMISS=1 AND (MSRF2= 'F' OR MSRF2='') then
ANTICOAGCODE61=.T

Else ANTICOAGCODE61=.

Source variable(s): FOUNDACOAG61, ALLMISS, MSRF2

8.7 ASPIRINCODE61 (V6 Aspirin Use in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: (From UC4892 – V4)
If FOUNDASP61=1 then ASPIRINCODE61=1

Else if FOUNDASP61=0 AND ((ALLMISS=1 AND MSRF2= 'T') OR
(ALLMISS=0)) then ASPIRINCODE61=0

Else if ALLMISS=1 AND (MSRF2= 'F' OR MSRF2='') then
ASPIRINCODE61=.T

Else ASPIRINCODE61=.

Source variable(s): FOUNDASP61, ALLMISS, MSRF2

8.8 **ANTIANXMDCODE61 (V6 Antianxiety Medication in past 4 weeks - Using Medi-Span GPI Code)**

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: (From UC4735 – V4)
If FOUNDANTIANX61=1 then ANTIANXMDCODE61=1

Else if FOUNDANTIANX61=0 AND ((ALLMISS=1 AND MSRF2= 'T')
OR (ALLMISS=0)) then ANTIANXMDCODE61=0

Else if ALLMISS=1 AND (MSRF2= 'F' OR MSRF2='') then
ANTIANXMDCODE61=.T

Else ANTIANXMDCODE61=.

Source variable(s): FOUNDANTIANX61, ALLMISS, MSRF2

8.9 **ANTIPSYCHMDCODE61 (V6 Antipsychotic Medication in past 4 weeks - Using Medi-Span GPI Code)**

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: (From UC4735 – V4)
If FOUNDANTIPSYCH61=1 then ANTIPSYCHMDCODE61=1

Else if FOUNDANTIPSYCH61=0 AND ((ALLMISS=1 AND
MSRF2='T') OR (ALLMISS=0)) then ANTIPSYCHMDCODE61=0

Else if ALLMISS=1 AND (MSRF2= 'F' OR MSRF2='') then
ANTIPSYCHMDCODE61=.T

Else ANTIPSYCHMDCODE61=.

Source variable(s): FOUNDANTIPSYCH61, ALLMISS, MSRF2

8.10 HYPNOTMDCODE61 (V6 Hypnotic/Sedative Medication in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: (From UC4735 – V4)
If FOUNDHYPNOT61=1 then HYPNOTMDCODE61=1

Else if FOUNDHYPNOT61=0 AND ((ALLMISS=1 AND MSRF2= 'T')
OR (ALLMISS=0)) then HYPNOTMDCODE61=0

Else if ALLMISS=1 AND (MSRF2= 'F' OR MSRF2='') then
HYPNOTMDCODE61=.T

Else HYPNOTMDCODE61=.

Source variable(s): FOUNDHYPNOT61, ALLMISS, MSRF2

8.11 ANTICONVMDCODE61 (V6 Anticonvulsant Medication in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: (From UC4735 – V4)
If FOUNDANTICONV61=1 then ANTICONVMDCODE61=1

Else if FOUNDANTICONV61=0 AND ((ALLMISS=1 AND MSRF2= 'T')
OR (ALLMISS=0)) then ANTICONVMDCODE61=0

Else if ALLMISS=1 AND (MSRF2= 'F' OR MSRF2='') then
ANTICONVMDCODE61=.T

Else ANTICONVMDCODE61=.

Source variable(s): FOUNDANTICONV61, ALLMISS, MSRF2

8.12 ANTIDEMMDCODE61 (V6 Antidementia/Nootropic Medication in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: (From UC4735 – V4)
If FOUNDANTIDEM61=1 then ANTIDEMMDCODE61=1

Else if FOUNDANTIDEM61=0 AND ((ALLMISS=1 AND MSRF2= 'T')
OR (ALLMISS=0)) then ANTIDEMMDCODE61=0

Else if ALLMISS=1 AND (MSRF2= 'F' OR MSRF2='') then
ANTIDEMMDCODE61=.T

Else ANTIDEMMDCODE61=.

Source variable(s): FOUNDANTIDEM61, ALLMISS, MSRF2

8.13 CNSALTMDCODE61 (V6 CNS Altering Medication in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: (From UC4735 – V4)
If FOUNDCNSATL61=1 then CNSALTMDCODE61=1

Else if FOUNDCNSATL61=0 AND ((ALLMISS=1 AND MSRF2= 'T')
OR (ALLMISS=0)) then CNSALTMDCODE61=0

Else if ALLMISS=1 AND (MSRF2= 'F' OR MSRF2='') then
CNSALTMDCODE61=.T

Else CNSALTMDCODE61=.

Source variable(s): FOUNDCNSATL61, ALLMISS, MSRF2

8.14 DIABMDCODE61 (V6 Diabetic Medications in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If FOUNDDIAB61=1 then DIABMDCODE61=1

Else if FOUNDDIAB61=0 AND ((ALLMISS=1 AND MSRF2= 'T') OR (ALLMISS=0)) then DIABMDCODE61=0

Else if ALLMISS=1 AND (MSRF2= 'F' OR MSRF2='') then DIABMDCODE61=.T

Else DIABMDCODE61=.

Source variable(s): FOUNDDIAB61, ALLMISS, MSRF2

8.15 BETAMDCODE61 (V6 Beta-Blocker in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If FOUNDBETA61=1 then BETAMDCODE61=1

Else if FOUNDBETA61=0 AND ((ALLMISS=1 AND MSRF2='T') OR (ALLMISS=0)) then BETAMDCODE61=0

Else if ALLMISS=1 AND (MSRF2='F' OR MSRF2='') then BETAMDCODE61=.T

Else BETAMDCODE61=.

Source variable(s): FOUNDBETA61, ALLMISS, MSRF2

8.16 ANGINHMDCODE61 (V6 Angiotensin converting enzyme inhibitor in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If FOUNDANGINH61=1 then ANGINHMDCODE61=1

Else if FOUNDANGINH61=0 AND ((ALLMISS=1 AND MSRF2='T') OR (ALLMISS=0)) then ANGINHMDCODE61=0

Else if ALLMISS=1 AND (MSRF2='F' OR MSRF2='') then ANGINHMDCODE61=.T

Else ANGINHMDCODE61=.

Source variable(s): FOUNDANGINH61, ALLMISS, MSRF2

8.17 ANGIANTMDCODE61 (V6 Angiotensin II receptor antagonists in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If FOUNDANGANT61=1 then ANGIANTMDCODE61=1

Else if FOUNDANGANT61=0 AND ((ALLMISS=1 AND MSRF2='T') OR (ALLMISS=0)) then ANGIANTMDCODE61=0

Else if ALLMISS=1 AND (MSRF2='F' OR MSRF2='') then ANGIANTMDCODE61=.T

Else ANGIANTMDCODE61=.

Source variable(s): FOUNDANGANT61, ALLMISS, MSRF2

8.18 ALDANTMDCODE61 (V6 Aldosterone Antagonist in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If FOUNDALDANT61=1 then ALDANTMDCODE61=1

Else if FOUNDALDANT61=0 AND ((ALLMISS=1 AND MSRF2='T') OR (ALLMISS=0)) then ALDANTMDCODE61=0

Else if ALLMISS=1 AND (MSRF2='F' OR MSRF2='') then ALDANTMDCODE61=.T

Else ALDANTMDCODE61=.

Source variable(s): FOUNDALDANT61, ALLMISS, MSRF2

8.19 LOOPDIUMDCODE61 (V6 Loop Diuretic in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If FOUNDLOOPDIU61=1 then LOOPDIUMDCODE61=1

Else if FOUNDLOOPDIU61=0 AND ((ALLMISS=1 AND MSRF2='T') OR (ALLMISS=0)) then LOOPDIUMDCODE61=0

Else if ALLMISS=1 AND (MSRF2='F' OR MSRF2='') then LOOPDIUMDCODE61=.T

Else LOOPDIUMDCODE61=.

Source variable(s): FOUNDLOOPDIU61, ALLMISS, MSRF2

8.20 DIGMDCODE61 (V6 Digoxin in past 4 weeks - Using Medi-Span GPI Code)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If FOUNDDIG61=1 then DIGMDCODE61=1

Else if FOUNDDIG61=0 AND ((ALLMISS=1 AND MSRF2='T') OR (ALLMISS=0)) then DIGMDCODE61=0

Else if ALLMISS=1 AND (MSRF2='F' OR MSRF2='') then DIGMDCODE61=.T

Else DIGMDCODE61=.

Source variable(s): FOUNDDIG61, ALLMISS, MSRF2

9. PHYSICAL ACTIVITY

9.1 SPRT_I61 (V6 Sport during Leisure Time)

Description: This index is a composite score of sport during leisure time.

Format: Continuous variable between 1 and 5

Type: Numeric

- Create temporary variables I2_1, I2_2, I2_3, I2_4 (sport score corresponding to each of the 4 recorded sports or exercises reported by the participant)

PAC variables being used in temporary variable					
	Sport Count	Activity	Time	Proportion	Remark
I2_1	1	PAC2	PAC3	PAC4	
I2_2	2	PAC6	PAC7	PAC8	0 if PAC5='N'
I2_3	3	PAC10	PAC11	PAC12	0 if PAC9='N'
I2_4	4	PAC14	PAC15	PAC16	0 if PAC13='N'

- An intensity level is assigned to each of the activity codes

Intensity	Activity Codes (PAC2, PAC6, PAC10, PAC14)
Light	1,31,43,55,60,61,73,76,79,97,121,125,136,142,154,169,178,208,229,244,286,304,322,328,403,412,415,418,421,448,499
Moderate	2,3,4,7,10,16,25,28,37,40,49,52,70,82,94,100,118,124,130,139,145,146,148,160,163,175,181,187,190,193,199,205,211,217,232,235,238,247,249,350,333,498,259,262,265,271,289,292,295,301,310,313,319,325,331,340,352,355,358,364,376,385,388,391,397,400,404,406,409,427,430,436,451
Heavy	13,19,22,46,58,67,85,88,91,109,112,115,127,133,151,157,166,172,184,196,202,214,220,223,226,241,250,253,268,274,277,280,283,298,316,334,337,343,346,349,361,367,370,373,379,382,394,424,433,437,439,442

- Calculate sport score for each activity using the following formula, intensity(X)*hours_weeks(Y)*months_year(Z) where X, Y, and Z are assigned values according to the following:

Intensity	X
Light	0.76
Moderate	1.26
Heavy	1.76

Hours (per week)	Variable	Y

	value	
Less than one hour	A	.5
At least 1 but not quite 2	B	1.5
At least 2 but not quite 3	C	2.5
At least 3 but not quite 4	D	3.5
4 or more	E	4.5

Months (per year)	Variable value	Z
Less than 1	A	0.04
At least 1 but not quite 4	B	0.17
At least 4 but not quite 7	C	0.42
At least 7 but not quite 10	D	0.67
10 or more	E	0.92

I2 Summary sports score

- Calculate $I_2 = I_{2_1} + I_{2_2} + I_{2_3} + I_{2_4}$: sum of four simple sports scores.

Note:

If PAC5='N' then $I_{2_2} = 0$

If PAC9='N' then $I_{2_3} = 0$

If PAC13='N' then $I_{2_4} = 0$

This sum score is then recoded to a score of 1 to 5 based on the following criteria:

If PAC1='N' then $I_2 = 1$ else

If $0 \leq I_2 < 0.01$ then $I_2 = 1$

If $0.01 \leq I_2 < 4$ then $I_2 = 2$

If $4 \leq I_2 < 8$ then $I_2 = 3$

If $8 \leq I_2 < 12$ then $I_2 = 4$

If $12 \leq I_2$ then $I_2 = 5$

I3 Leisure sport exercise activity versus peers, recoded

- PAC19 was recoded as follows:

Format	PAC19 value	Recoded value for algorithm
Much less	A	1
Less	B	2
The same	C	3
More	D	4
Much more	E	5

I4 Sweat during leisure time, recoded

- PAC20 was recoded as follows:

Format	PAC20 value	Recoded value for algorithm
Never	A	1
Seldom	B	2
Sometimes	C	3
Often	D	4
Very often	E	5

I5 Sports/exercise during leisure activity, recoded

- PAC18 was recoded as follows:

Format	PAC18 value	Recoded value for algorithm
Never	A	1
Seldom	B	2
Sometimes	C	3
Often	D	4
Very often	E	5

$SPRT_I61 = (I2+I3+I4+I5)/4$

If I2 or I3 or I4 or I5 is missing, then $SPRT_I61 = .T$

9.2 LISR_I62 (V6 Physical Activity during Leisure Time Excluding Sport)

Description: This index is a composite score of leisure time activity that includes frequency of TV viewing, frequency of walking, and frequency of bicycling. The question about the number of minutes walked or bicycled per day to and from work or shopping was not included in the battery so the variable is not exactly defined as was in previous visits.

Format: Continuous variable between 1 and 5

Type: Numeric

I6 Leisure time television watching, recoded

- PAC21 was recoded as follows:

Format	PAC21 value	Recoded value for algorithm
Never	A	5
Seldom	B	4
Sometimes	C	3
Often	D	2
Very often	E	1

I7 Leisure time walking frequency, recoded

- PAC22 was recoded as follows:

Format	PAC22	Recoded value for algorithm
Never	A	1
Seldom	B	2
Sometimes	C	3
Often	D	4
Very often	E	5

I8 Leisure time cycling frequency, recoded

- PAC23 was recoded as follows:

Format	PAC23 value	Recoded value for algorithm
Never	A	1
Seldom	B	2
Sometimes	C	3
Often	D	4
Very often	E	5

$LISR_I62 = (I6 + I7 + I8 + ((I7 + I8) / 2)) / 4$

If I6 or I7 or I8 is missing, then $LISR_I62 = .T$

10. PHYSICAL FUNCTION

The physical function variables mainly use the data collected on the ARIC PFX form. The short physical performance battery (SPPB) is a group of measures that combines the results of the gait speed, chair stand and balance tests (Guralnik et al., 2000). The scores range from 0 (worst performance) to 12 (best performance). Frailty variables, originally derived at Visit 5, are also described in this section.

10.1 SPPBCS61 (V6 Physical Function Chair Stand)

Format: integer variable (0-4 possible points)

Type: Numeric

Algorithm: If (PFX1 in (2,3,4)) OR (PFX2 in (2,3)) OR (PFX2b_DER \geq 60) then SPPBCS61=0;
Else if (16.70 \leq PFX2b_DER <60) then SPPBCS61=1
Else if (13.70 \leq PFX2b_DER <16.70) then SPPBCS61=2
Else if (11.20 \leq PFX2b_DER <13.70) then SPPBCS61=3
Else if (. <PFX2b_DER <11.20) then SPPBCS61=4
Else SPPBCS61=.

Source variable(s): PFX1, PFX2, PFX2b_DER

10.2 SPPBST61 (V6 Physical Function Semi Tandem Stand)

Format: integer variable

Type: Numeric

Algorithm: If (PFX3 in (2,3,4)) then SPPBST61=0
Else if (PFX3=5) then SPPBST61=1
Else SPPBST61=.

Source variable(s): PFX3

10.3 SPPBSBS61 (V6 Physical Function Side-by-Side Stand)

Format: integer variable

Type: Numeric

Algorithm: If (PFX4 in (2,3,4)) then SPPBSBS61=0;
Else if (PFX3=5 OR PFX4=5) then SPPBSBS61=1;
Else SPPBSBS61=.

Source variable(s): PFX3, PFX4

10.4 SPPBTS61 (V6 Physical Function Tandem Stand)

Format: integer variable

Type: Numeric

Algorithm: If max(PFX5, PFX6) in (2,3,4) then SPPBTS61=0;
else if $3 \leq \max(\text{PFX5a_DER}, \text{PFX6a_DER}) < 10$ then SPPBTS61=1;
else if max(PFX5, PFX6) in (5) then SPPBTS61 = 2;
else SPPBTS61=.;

Source variable(s): PFX5, PFX5a_DER, PFX6, PFX6a_DER

10.5 SPPBBAL61 (V6 Physical Function Summary Balance Score)

Format: integer variable

Type: Numeric

Algorithm: SPPBBAL61=sum(of SPPBST61,SPPBSBS61,SPPBTS61);

Source variable(s): SPPBSBS61, SPPBST61, SPPBTS61

10.6 WALK4M61 (V6 Physical Function 4 Meter Walk, Fastest Time of 2 Trials)

Format: continuous variable (f4.2)

Type: Numeric

Algorithm: The better of (1) PFX7a and PFX8a if walking aid is used OR the better of (2) PFX7b and PFX8b if no walking aid is used. There are some instances where a walking aid is used in one trial and not in the other trial. The variable will be the fastest time walked regardless of using the aid or not. All participants at the clinic visits should have a timed walk. Any who did not do the test due to "not attempted/unable" will have a missing value for this variable. If only one trial completed, use the results from that trial.

If $\max(\text{pfx7}, \text{pfx8}) \leq 2$ then WALK4M61=.T;

Else
WALK4M61=min(PFX7a_der,PFX7b_der,PFX8a_der,PFX8b_der);

Source variable(s): PFX7, PFX7a_der, PFX7b_der, PFX8, PFX8a_der, PFX8b_der

10.7 WALKAID61 (V6 Physical Function 4 Meter Walk: Used Walking Aid)

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: This indicator variable will be set to 1 when a patient used a walking aid during the 4 meter walk. The walk is performed twice and in the instance where a patient uses an aid in one trial, but not in the other, this variable will be set according to the presence or absence of a walking aid that goes with the fastest time of the 2 trials.

If WALK4M61>.T and (WALK4M61=PFX7a_der or
WALK4M61=PFX8a_der) then WALKAID61 = 1;
Else if WALK4M61>.T then WALKAID61 = 0;
Else WALKAID61=.T;

Source variable(s): PFX7a_der, PFX8a_der, WALK4M61

10.8 WALK4M62 (V6 Physical Function 4 Meter Walk, Average Time of 2 Trials (BOTH TRIAL WITH AID OR BOTH TRIALS WITHOUT AID))

Format: continuous variable (f4.2)

Type: Numeric

Algorithm: WALK4M62= MAX((PFX7A_DER+PFX8A_DER)/2, (PFX7B_DER +
PFX8B_DER)/2)

Source variable(s): PFX7A_DER, PFX7B_DER, PFX8A_DER, PFX8B_DER

10.9 WALKAID62 (V6 Physical Function 4 Meter Walk: Used Walking Aid in both trials (USE WITH WALK4M62))

Format: 0=No,
1=Yes,
.T=missing.

Type: Numeric

Algorithm: If WALK4M62>.T and (PFX7=3 and PFX8=3) then WALKAID62 = 1;
Else if WALK4M62>.T then WALKAID62=0;
Else WALKAID62=.T;

Source variable(s): PFX7, PFX8, WALK4M62

10.10 SPPB4M61 (V6 Physical Function 4 Meter Walk Score)

Format: integer variable (0-4)

Type: Numeric

Algorithm: If max(PFX7,PFX8) = 2 then SPPB4M61 = 0;
Else if (WALK4M61 > 8.70) SPPB4M61 = 1;
Else if (6.21 <= WALK4M61 <= 8.70) then SPPB4M61 = 2;
Else if (4.82 <= WALK4M61 < 6.21) then SPPB4M61 = 3;
Else if (.z < WALK4M61 < 4.82) then SPPB4M61 = 4;
Else SPPB4M61 = .T ;

Source variable(s): PFX7, PFX8, WALK4M61

10.11 SPPB61 (V6 Short Physical Performance Summary Battery Score)

Format: integer variable

Type: Numeric

Algorithm: SPPB61=sum(of SPPBCS61,SPPBBAL61,SPPB4M61);

Source variable(s): SPPB4M61, SPPBBAL61, SPPBCS61

10.12 GRIPBEST61 (V6 Physical Function Grip, Best of 2 Trials)

Format: continuous variable

Type: Numeric

Algorithm: Max of PFX11b and PFX11c. Only participants who respond "Both" to PFX10a should be excluded (i.e. surgery on both hands).

If PFX10 NE missing and PFX10a NE 'B' then
GRIPBEST61=max(PFX11b,PFX11c);

Source variable(s): PFX10a, PFX11b, PFX11c

10.13 GRIPMEAN61 (V6 Physical Function Grip, Mean of 2 Trials or Result for 1 Trial)

Format: continuous variable

Type: Numeric

Algorithm: Mean of PFX11b and PFX11c. Only participants who respond "Both" to PFX10a should be excluded (i.e. surgery on both hands).

If PFX10 NE missing and PFX10a NE 'B' then
GRIPMEAN61=mean(PFX11b,PFX11c);

Source variable(s): PFX10a, PFX11b, PFX11c

10.14 V5V6WTDELTA61 (% of V5 weight change from V5 to V6 (neg val means loss) – frailty weight loss component)

Description: The change in weight from visit 5 to visit 6 is calculated and presented as the % of visit 5 weight.

Format: continuous variable (0-99, %)

Type: Numeric

Algorithm: $((ant4-v5ant4)/v5ant4)*100$, where v5ant4 = ant4 from visit 5 and ant4=ant4 from visit 6

Source variable(s): ANT4, V5ANT4

10.15 WALKSPEED15FT61 (Time in seconds used to walk 15ft – frailty slowness component)

Description: The physical function form collects data on times to walk 4 meters. The WALKSPEED15FT61 variable uses the 4m walk rate to calculate the time in seconds needed to walk 15ft.

Format: numeric continuous variable

Type: Numeric

Algorithm: WALKSPEED15FT61=MIN(((15/3.28084)* (pfx7a_der/4), (15/3.28084)*(pfx7b_der/4), (15/3.28084)*(pfx8a_der/4), (15/3.28084)*(pfx8b_der/4))

Source variable(s): PFX7A_DER, PFX7B_DER, PFX8A_DER, PFX8B_DER

10.16 V6CANCER61 (Reported a cancer site during an ARIC follow-up interview in the time from 06-15-2016 to the V6date61 (V6 visit date))

Description: Indicator used as proxy measure for having cancer at V6.

Format: 0=No,
1=Yes,
.=missing.

Type: Numeric

Algorithm: The V6CANCER61 is a proxy measure derived from ARIC follow-up data. If a PPT reported a cancer site during their follow-up interview that occurred anytime between June 15, 2016 and the PPT visit date then the variable was set to 1. If the PPT was interviewed for ARIC follow-up during that time and did not report a cancer site then the variable was set to 0, otherwise the variable was set to missing.

Source variable(s):

10.17 TR1WALK4MSP61 (V6 4M Walking Speed for Trial 1 (m/sec))

Format: continuous variable

Type: Numeric

Algorithm: TR1WALK4MSP61=4/(max(PFX7a_der, PFX7b_der)

Source variable(s): PFX7a, PFX7b

10.18 TR2WALK4MSP61 (V6 4M Walking Speed for Trial 2 (m/sec))

Format: continuous variable

Type: Numeric

Algorithm: TR2WALK4MSP61=4/(max(PFX8a_der, PFX8b_der)

Source variable(s): PFX8a, PFX8b

10.19 AVGWALK4MSP61 (V6 Average 4M Walking Speed (m/sec))

Format: continuous variable

Type: Numeric

Algorithm: $AVGWALK4MSP61 = \text{mean}(TR1WALK4MSP61, TR2WALK4MSP61)$

Source variable(s): TR1WALK4MSP61, TR2WALK4MSP61

10.20 MINWALK4MSP61 (V6 Fastest 4M Walking Speed for Both Trials (m/sec))

Format: continuous variable

Type: Numeric

Algorithm: $MINWALK4MSP61 = \text{min}(TR1WALK4MSP61, TR2WALK4MSP61)$;

Source variable(s): TR1WALK4MSP61, TR2WALK4MSP61

10.21 UNABLETOWALK61 (Indicator Variable Noting PPT No Attempt to Walk 4M, Not Able)

Format: 0=No,
1=Yes,
.=Missing.

Type: Numeric

Algorithm: If PFX7 and PFX8 are both NULL then UNABLETOWALK61=NULL
Else if PFX7=2 and PFX8 = missing or 2 then UNABLETOWALK61=1
Else UNABLETOWALK61=0

Source variable(s): PFX7, PFX8

10.22 UNINTEND_WTLOSS61 (V6 Unintentional Weight Loss)

Format: 0=No,
1=Yes,
.=missing.

Type: Numeric

Algorithm: If tmw1=1 and tmw3=0 then UNINTEND_WTLOSS61=1;
Else if tmw1=1 then UNINTEND_WTLOSS61=0;

Source variable(s): TMW1, TMW3

10.23 TMW_TOTFT61 (V6 Total Feet Walked in 2-Minute Walk)

Format: continuous variable

Type: Numeric

Algorithm: $TMW_TOTFT61 = \text{sum}((tmw8 * 50), tmw9);$

Source variable(s): TMW8, TMW9

10.24 TMW_COMPSPEEDFTPERSEC61 (V6 TMW Speed for Completers (ft/sec))

Format: continuous variable

Type: Numeric

Algorithm: Calculate if tmw_pres=1 and tmw11=5
 $TMW_COMPSPEEDFTPERSEC61 = tmw_totft61 / 120;$

Source variable(s): TMW11, TMW_TOTFT61

10.25 TMW_COMPSPEEDMPERSEC61 (V6 TMW Speed for Completers (m/sec))

Format: continuous variable

Type: Numeric

Algorithm: Calculate if tmw_pres=1 and tmw11=5
 $TMW_COMPSPEEDMPERSEC61 = tmw_totft61 * 0.3048 / 120$

Source variable(s): TMW11, TMW_TOTFT61

10.26 TMW_NONCOMPSECWALK61 (V6 TMW Seconds Walking for Noncompleters)

Format: continuous variable

Type: Numeric

Algorithm: Calculate if tmw_pres=1 and tmw11=4
 $TMW_NONCOMPSECWALK61 = \text{sum}((tmw12a*60), tmw12b)$

Source variable(s): TMW11, TMW12a, TMW12b

10.27 TMW_NONCOMPSPEEDFTPERSEC61 (V6 TMW Speed for Noncompleters (ft/sec))

Format: continuous variable

Type: Numeric

Algorithm: Calculate if tmw_pres=1 and tmw11=4 and
 $tmw_noncompsecwalk61 > 0$
 $TMW_NONCOMPSPEEDFTPERSEC61 = tmw_totft61 / tmw_noncompsecwalk61$

Source variable(s): tmw11, tmw_noncompsecwalk61, tmw_totft61

10.28 TMW_NONCOMPSPEEDMPERSEC61 (V6 TMW Speed for Noncompleters (m/sec))

Format: continuous variable

Type: Numeric

Algorithm: Calculate if tmw_pres=1 and tmw11=4 and
 $tmw_noncompsecwalk61 > 0$
 $TMW_NONCOMPSPEEDMPERSEC61 = tmw_totft61 * 0.3048 / tmw_noncompsecwalk61$

Source variable(s): tmw11, tmw_noncompsecwalk61, tmw_totft61

10.29 PAQ_LIMITEDFUNCTION61 (V6 PAQ Limited Function)

Description: Indicator variable that is set to 1 (Yes) when the PPT indicates they need assistance for items listed on the Physical Abilities Questionnaire (PAQ).

Format: 0=No
1=Yes
.=Missing

Type: Numeric

Algorithm: PAQ_LIMITEDFUNCTION61 = (('B'<=paq1<='D') OR =
('B'<=paq2<='D') OR ('B'<=paq3<='D') OR ('B'<=paq4<='D') OR
('B'<=paq5<='D') OR ('B'<=paq6<='D') OR ('B'<=paq7<='D') OR
('B'<=paq8<='D') OR ('B'<=paq9<='D') OR ('B'<=paq10<='D') OR
('B'<=paq11<='D') OR ('B'<=paq12<='D') OR (paq13=1) OR
(paq14=1) OR (paq15=1) OR (paq16=1) OR (paq17=1))

Source variable(s): PAQ1, PAQ2, PAQ3, PAQ4, PAQ5, PAQ6, PAQ7, PAQ8, PAQ9,
PAQ10, paq11, paq12, paq13, paq14, paq15, paq16, paq17

10.30 PAQ_GOODFUNCTION61 (V6 PAQ Good Function)

Description: Indicator variable that is set to 1 (Yes) when the PPT indicates they need NO assistance for items listed on the Physical Abilities Questionnaire (PAQ).

Format: 0=No
1=Yes
.=Missing

Type: Numeric

Algorithm: PAQ_GOODFUNCTION61= ((paq1='A') AND (paq2='A') AND
(paq3='A') AND (paq4='A') AND (paq5='A') AND (paq6='A') AND
(paq7='A') AND (paq8='A') AND (paq9='A') AND (paq10='A') AND
(paq11='A') AND (paq12='A') AND (paq13=0) AND (paq14=0) AND
(paq15=0) AND (paq16=0) AND (paq17=0))

Source variable(s): PAQ1, PAQ2, PAQ3, PAQ4, PAQ5, PAQ6, PAQ7, PAQ8, PAQ9,
PAQ10, PAQ11, PAQ12, PAQ13, PAQ14, PAQ15, PAQ16, PAQ17

10.31 EXHAUST61 (V6 Responded 2 or 3 on CES3 or CES11 (CESD) – frailty exhaustion component)

Description: Second component in considering frailty of PPT based on exhaustion from depression.

Format: 0=No
1=Yes
.=Missing

Type: Numeric

Algorithm: EXHAUST61=1, if (CES3=2 OR CES3=3) OR (CES11=2 OR CES11=3)
Else EXHAUST61= NULL, if CES3= NULL AND CES11= NULL
ELSE EXHAUST61=0

Source variable(s): CES2, CES3, CES11

10.32 LOWENERGYCOMP61 (V6 Lowest quintile of SPRT_i61 - frailty low physical activity component)

Description: The third component in considering frailty of PPT based on low physical activity during leisure time.

Format: 0=No
1=Yes
.=Missing

Type: Numeric

Algorithm: LOWENERGYCOMP61=1, if (<SPRT_I61<1.75 AND GENDER='F')
OR (SPRT_I61<2.00 AND GENDER='M')

LOWENERGYCOMP61=0, if (<SPRT_I61>=1.75 AND GENDER='F')
OR (SPRT_I61>=2.00 AND GENDER='M')

LOWENERGYCOMP61=NULL, if gender=NULL or SPRT_I61=NULL

Source variable(s): SPRT_I61, gender

10.33 WTLOSSCOMPA61 (V6 Lost >5% weight or Low BMI (<18.5) – frailty weight loss component for 61a definition)

Description: First component in considering frailty of PPT based on weight loss greater than 5% or low BMI.

Format: 0=No
1=Yes
.=Missing

Type: Numeric

Algorithm: WTLOSSCOMP61=1, if (<V5V6WTDELTAP61<=-5) OR (<BMI61<18.5)

WTLOSSCOMP61=0, if (V5V6WTDELTA61>-5) AND (BMI61>=18.5)

WTLOSSCOMP61=NULL, if V5V6WTDELTA61 and BMI61 are NULL

Source variable(s): V5V6WTDELTA61, BMI61

10.34 WTLOSSCOMP61 (V6 Lost >10% weight or Low BMI (<18.5) – frailty weight loss component for 61b definition)

Description: First component in considering frailty of PPT based on more restrictive weight loss greater than 10% or low BMI.

Format:
0=No
1=Yes
.=Missing

Type: Numeric

Algorithm: WTLOSSCOMP61=1, if (.<V5V6WTDELTA61<=-10) OR (.<BMI61<18.5)

WTLOSSCOMP61=0, if (V5V6WTDELTA61>-10) AND (BMI61>=18.5)

WTLOSSCOMP61=NULL, if V5V6WTDELTA61 and BMI61 are NULL

Source variable(s): V5V6WTDELTA61, BMI61

10.35 WALKSPEEDCOMP61 (V6 Slowest 20% time to walk 15ft – frailty slowness component)

Description: Fourth component in considering frailty of PPT based on slowness by walking.

Format:
0=No
1=Yes
.=Missing

Type: Numeric

Algorithm: WALKSPEEDCOMP61=1:

Males:

IF (gender = 'M' AND V6IN129 <=173 AND Walkspeed15ft61>= 7) OR
IF (gender = 'M' AND V6IN129 > 173 AND Walkspeed15ft61>= 6) OR

Females:

IF (gender = 'F' AND V6IN129 <=159 AND Walkspeed15ft61>= 7) OR
IF (gender = 'F' AND V6IN129 > 159 AND Walkspeed15ft61>= 6)

WALKSPEEDCOMP61=NULL:

IF gender=NULL or V6IN129=NULL or Walkspeed15ft61= NULL
ELSE WALKSPEEDCOMP61=0

Source variable(s): gender, V6IN129, Walkspeed15ft61

10.36 GRIPSTRENGTHCOMP61 (V6 Slowest 20% grip strength – frailty weakness component)

Description: Fifth component in considering frailty of PPT based on grip strength and BMI.

Format: 0=No
1=Yes
.=Missing

Type: Numeric

Algorithm: GRIPSTRENGTHCOMP61=1:

Males:

IF gender = 'M' AND BMI61 <= 24 AND .<MAX(pfx11b, pfx11c) <= 29
OR

IF gender = 'M' AND 24 < BMI61 <= 26 AND .<MAX(pfx11b, pfx11c)
<= 30 OR

IF gender = 'M' AND 26 < BMI61 <= 28 AND .<MAX(pfx11b, pfx11c)
<= 30 OR

IF gender = 'M' AND BMI61 > 28 AND .<MAX(pfx11b, pfx11c) <= 32
OR

Females:

IF gender = 'F' AND BMI61 <= 23 AND .<MAX(pfx11b, pfx11c) <= 17
OR

IF gender = 'F' AND 23 < BMI61 <= 26 AND .<MAX(pfx11b, pfx11c)
<= 17.3 OR

IF gender = 'F' AND 26 < BMI61 <= 29 AND .<MAX(pfx11b, pfx11c)
<= 18 OR

IF gender = 'F' AND BMI61 > 29 AND .<MAX(pfx11b, pfx11c) <= 21

GRIPSTRENGTHCOMP61=NULL:
IF gender=NULL or BMI61=NULL or (pfx11b and pfx11c)=NULL

GRIPSTRENGTHCOMP61=0:
Otherwise

Source variable(s): gender, BMI61, pfx11b, pfx11c

10.37 FRAILTY61a (V6 ARIC Physical Function WG Frailty Definition (a))

Description: There are 5 components considered in deriving the frailty variable. Indicators for these components are defined below. If 3 or more of the components are present, then FRAILTY61a=1. If 1 or 2 of the components are present, then FRAILTY61a=2. Of none of the components are present and there is data for each component, then FRAILTY61a=3.

Format: 1=Frail
2=Prefrail
3=Robust
.=Missing

Type: Numeric

Algorithm:

1. Weight loss:
If (.<V5V6WTDELTA61<=-5) OR (.<BMI61<18.5)
2. Exhaustion:
If EXHAUST61=1
3. Low Energy:
If (.<SPRT_I61<1.75 AND GENDER='F') OR (.<SPRT_I61<2.00 AND GENDER='M')
4. Walking Speed:
IF (gender = 'M' AND ant3 <=173 AND Walkspeed15ft61>= 7) OR
IF (gender = 'M' AND ant3 > 173 AND Walkspeed15ft61>= 6) OR
IF (gender = 'F' AND ant3 <=159 AND Walkspeed15ft61>= 7) OR
IF (gender = 'F' AND ant3 > 159 AND Walkspeed15ft61>= 6)
5. Grip Strength
IF gender = 'M' AND BMI61 <= 24 AND .<MAX(pfx11b, pfx11c) <= 29
OR

IF gender = 'M' AND 24 < BMI61 <= 26 AND .<MAX(pfx11b, pfx11c) <= 30 OR
 IF gender = 'M' AND 26 < BMI61 <= 28 AND .<MAX(pfx11b, pfx11c) <= 30 OR
 IF gender = 'M' AND BMI61 > 28 AND .<MAX(pfx11b, pfx11c) <= 32 OR
 IF gender = 'F' AND BMI61 <= 23 AND .<MAX(pfx11b, pfx11c) <= 17 OR
 IF gender = 'F' AND 23 < BMI61 <= 26 AND .<MAX(pfx11b, pfx11c) <= 17.3 OR
 IF gender = 'F' AND 26 < BMI61 <= 29 AND .<MAX(pfx11b, pfx11c) <= 18 OR
 IF gender = 'F' AND BMI61 > 29 AND .<MAX(pfx11b, pfx11c) <= 21

Source variable(s): V5V6WTDELTA61, BMI61, EXHAUST61, SPRT_I61, GENDER, ANT3 WALKSPEED15FT61, PFX11B, PFX11C

10.38 FRAILTY61b (V6 ARIC Physical Function WG Frailty Definition (b))

Description: There are 5 components considered in deriving the frailty variable. Indicators for these components are defined below. If 3 or more of the components are present, then FRAILTY61b=1. If 1 or 2 of the components are present, then FRAILTY61b=2. Of none of the components are present and there is data for each component, then FRAILTY61b=3.

Format: 1=Frail
 2=Prefrail
 3=Robust
 .=Missing

Type: Numeric

Algorithm:

1. Weight loss:
 If (.<V5V6WTDELTA61<=-10) OR (.<BMI61<18.5)
2. Exhaustion:
 If EXHAUST61=1
3. Low Energy:
 If (.<SPRT_I61<1.75 AND GENDER='F') OR (.<SPRT_I61<2.00 AND GENDER='M')
4. Walking Speed:
 IF (gender = 'M' AND ant3 <=173 AND Walkspeed15ft61>= 7) OR

IF (gender = 'M' AND ant3 > 173 AND Walkspeed15ft61 >= 6) OR
IF (gender = 'F' AND ant3 <= 159 AND Walkspeed15ft61 >= 7) OR
IF (gender = 'F' AND ant3 > 159 AND Walkspeed15ft61 >= 6)

5. Grip Strength

IF gender = 'M' AND BMI61 <= 24 AND .<MAX(pfx11b, pfx11c) <= 29
OR
IF gender = 'M' AND 24 < BMI61 <= 26 AND .<MAX(pfx11b, pfx11c)
<= 30 OR
IF gender = 'M' AND 26 < BMI61 <= 28 AND .<MAX(pfx11b, pfx11c)
<= 30 OR
IF gender = 'M' AND BMI61 > 28 AND .<MAX(pfx11b, pfx11c) <= 32
OR
IF gender = 'F' AND BMI61 <= 23 AND .<MAX(pfx11b, pfx11c) <= 17
OR
IF gender = 'F' AND 23 < BMI61 <= 26 AND .<MAX(pfx11b, pfx11c)
<= 17.3 OR
IF gender = 'F' AND 26 < BMI61 <= 29 AND .<MAX(pfx11b, pfx11c)
<= 18 OR
IF gender = 'F' AND BMI61 > 29 AND .<MAX(pfx11b, pfx11c) <= 21

Source variable(s): V5V6WTDELTA61, BMI61, EXHAUST61, SPRT_I61, GENDER,
ANT3 WALKSPEED15FT61, PFX11B, PFX11C

**10.39 FRAILTY63 (V6 ARIC Physical Function WG Frailty Definition – weight loss
dropped)**

Description: There are 4 components considered in deriving the frailty variable. Indicators for these components are defined below. If 3 or more of the components are present, then frailty63=1. If 1 or 2 of the components are present, then frailty63=2. If none of the components are present (=0) AND there is data for each component (meaning no component has a missing value), then frailty63=3.

Format: 1=Frail
2=Prefrail
3=Robust
.=Missing

Type: Numeric

Algorithm:

1. Exhaustion:
If EXHAUST61=1
2. Low Energy:

If (.<SPRT_I61<1.75 AND GENDER='F') OR (.<SPRT_I61<2.00 AND GENDER='M')

3. Walking Speed:

IF (gender = 'M' AND ant3 <=173 AND Walkspeed15ft61>= 7) OR
IF (gender = 'M' AND ant3 > 173 AND Walkspeed15ft61>= 6) OR
IF (gender = 'F' AND ant3 <=159 AND Walkspeed15ft61>= 7) OR
IF (gender = 'F' AND ant3 > 159 AND Walkspeed15ft61>= 6)

4. Grip Strength

IF gender = 'M' AND BMI61 <= 24 AND .<MAX(pfx11b, pfx11c) <= 29
OR
IF gender = 'M' AND 24 < BMI61 <= 26 AND .<MAX(pfx11b, pfx11c)
<= 30 OR
IF gender = 'M' AND 26 < BMI61 <= 28 AND .<MAX(pfx11b, pfx11c)
<= 30 OR
IF gender = 'M' AND BMI61 > 28 AND .<MAX(pfx11b, pfx11c) <= 32
OR
IF gender = 'F' AND BMI61 <= 23 AND .<MAX(pfx11b, pfx11c) <= 17
OR
IF gender = 'F' AND 23 < BMI61 <= 26 AND .<MAX(pfx11b, pfx11c)
<= 17.3 OR
IF gender = 'F' AND 26 < BMI61 <= 29 AND .<MAX(pfx11b, pfx11c)
<= 18 OR
IF gender = 'F' AND BMI61 > 29 AND .<MAX(pfx11b, pfx11c) <= 21

Source variable(s): EXHAUST61, SPRT_I61, GENDER, ANT3 WALKSPEED15FT61, PFX11B, PFX11C

10.40 FRAILTY64 (V6 ARIC Physical Function WG Frailty Definition (unintentional weight loss))

Description: There are 5 components considered in deriving the frailty variable. Indicators for these components are defined below. If 3 or more of the components are present, then frailty64=1. If 1 or 2 of the components are present, then frailty64=2. If none of the components are present AND there is data for each component, then frailty64=3.

Format: 1=Frail
2=Prefrail
3=Robust
.=Missing

Type: Numeric

Algorithm:

1. Unintentional weight loss
If UNINTEND_WTLOSS=1
2. Exhaustion:
If EXHAUST61=1
3. Low Energy:
If (.<SPRT_I61<1.75 AND GENDER='F') OR (.<SPRT_I61<2.00 AND GENDER='M')
4. Walking Speed:
IF (gender = 'M' AND ant3 <=173 AND Walkspeed15ft61>= 7) OR
IF (gender = 'M' AND ant3 > 173 AND Walkspeed15ft61>= 6) OR
IF (gender = 'F' AND ant3 <=159 AND Walkspeed15ft61>= 7) OR
IF (gender = 'F' AND ant3 > 159 AND Walkspeed15ft61>= 6)
5. Grip Strength
IF gender = 'M' AND BMI61 <= 24 AND .<MAX(pfx11b, pfx11c) <= 29
OR
IF gender = 'M' AND 24 < BMI61 <= 26 AND .<MAX(pfx11b, pfx11c)
<= 30 OR
IF gender = 'M' AND 26 < BMI61 <= 28 AND .<MAX(pfx11b, pfx11c)
<= 30 OR
IF gender = 'M' AND BMI61 > 28 AND .<MAX(pfx11b, pfx11c) <= 32
OR
IF gender = 'F' AND BMI61 <= 23 AND .<MAX(pfx11b, pfx11c) <= 17
OR
IF gender = 'F' AND 23 < BMI61 <= 26 AND .<MAX(pfx11b, pfx11c)
<= 17.3 OR
IF gender = 'F' AND 26 < BMI61 <= 29 AND .<MAX(pfx11b, pfx11c)
<= 18 OR
IF gender = 'F' AND BMI61 > 29 AND .<MAX(pfx11b, pfx11c) <= 21

Source variable(s): UNINTEND_WTLOSS61, EXHAUST61, SPRT_I61, GENDER, ANT3
WALKSPEED15FT61, PFX11B, PFX11C

10.41 GAITSPEED61 (V6 Gait Speed (m/sec))

Type: Numeric

Algorithm: If WALK4M61>0 then GAITSPEED61=4/WALK4M61

Source variable(s): WALK4M61

11. HEARING RESULTS

11.1 PTARIGHT61 (Four Frequency Pure-tone Average (PTA) of the Right Ear (Db HL))

Format: numeric continuous variable

Type: Numeric

Algorithm: If AUD4a1 and AUD4a7 are both not missing and the absolute value of (AUD4a1 - AUD4a7) > 10 then PTARIGHT61=NULL
Else if AUD0c="H" and AUD3="N" then PTARIGHT61=NULL
Else if AUD4a3 and AUD4a1 and AUD4a9 and AUD4a13 are all not missing then
PTARIGHT61=mean(AUD4a3,AUD4a1,AUD4a9,AUD4a13)

Source variable(s): AUD0c, AUD3, AUD4a1, AUD4a3, AUD4a7, AUD4a9, AUD4a13

11.2 PTALEFT61 (Four Frequency Pure-tone Average (PTA) of the Left Ear (Db HL))

Format: continuous variable

Type: Numeric

Algorithm: If AUD4b1 and AUD4b7 are both not missing and the absolute value of (AUD4b1 - AUD4b7) > 10 then PTALEFT61=NULL
Else if AUD0c="H" and AUD3="N" then PTALEFT61=NULL
Else if AUD4b3 and AUD4b1 and AUD4b9 and AUD4b13 are all not missing then
PTALEFT61=mean(AUD4b3,AUD4b1,AUD4b9,AUD4b13)

Source variable(s): AUD0c, AUD3, AUD4b1, AUD4b3, AUD4b7, AUD4b9, AUD4b13

11.3 PTABETTER61 (Four Frequency Pure-tone Average (PTA) of the Better Ear (smaller PTA) (Db HL))

Format: continuous integer

Type: Numeric

Algorithm: If PTARIGHT61 not missing or PTALEFT61 not missing then
PTABETTER61=min(PTARIGHT61, PTALEFT61)

Source variable(s): PTARIGHT61, PTALEFT61

11.4 PTABETTERCAT61 (Hearing Loss Category Based on the Four Frequency Pure-tone Average (PTA) of the B)

Description: Values of this categorical variable indicate the hearing loss category for hearing in the better ear. This variable is used in the PPT's summary of results.

Format: 1=No hearing loss,
2=Mild hearing loss,
3=Moderate hearing loss,
4=Severe hearing loss

Type: Numeric

Algorithm: If NULL < PTABETTER61 <= 25 then PTABETTER61=1
Else if 25 < PTABETTER61 <= 40 then PTABETTER61=2
Else if 40 < PTABETTER61 <= 70 then PTABETTER61=3
Else if PTABETTER61 > 70 then PTABETTER61=4

Source variable(s): PTABETTER61

12. NEUROCOGNITIVE STUDY

12.1 CESD61 (V6 CES-Depression Scale)

Format: Continuous Numeric Variable

Type: numeric

Algorithm: If there are 10 non-missing items among CES1 through CES11 then CESD61=sum of CES1 through CES11
Otherwise missing.

Source variable(s): CES1 – CES11

12.2 FAQ61 (Functional Activities Questionnaire)

Description: Numeric variable for score on the Functional Activities Questionnaire.

Type: Numeric

Algorithm: IF [V6NCSCOG61] SELECTEDSTG2= "Y" then CALCULATE:
FAQ61=CDI25 + CDI26 + CDI31 + 2(CDI35) + CDI36 + CDI37 +
CDI18 + CDI17 + CDI22

Source variable(s): CDI17, CDI18, CDI22, CDI25, CDI26, CDI31, CDI35, CDI36, CDI37

13. SF-12 OUTCOMES

The proposed variables are based on the SF-12 questionnaire that was administered as part of Visit 6 using form SFE. All questions in that form will be used to derive the SF12 variables. Those missing a response on any question that is part of the SF12 scale item will be missing that scale, but not other scales for which they have non-missing responses. Those missing individual scales will have missing summary scores for which the missing scale is a component.

The company that holds the license has instructed us to include the statement "a modified version of the SF-12v2 was used" in ALL manuscripts using SF-12 data.

Note that the modifications are very minor, things like underlining certain words or phrases in the questions, and largely irrelevant because the instrument was interviewer-administered.

13.1 SF12PF61 (V6 SF-12 Physical Functioning Scale)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: $SF12PF61=100*(((SFE2a+SFE2b)-2)/4);$

Source variable(s): SFE2a, SFE2b

13.2 SF12RP61 (V6 SF-12 Role Physical Scale)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: $SF12RP61=100*(((SFE3a+SFE3b)-2)/8);$

Source variable(s): SFE3a, SFE3b

13.3 SF12BP61 (V6 SF-12 Bodily Pain Scale)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Note: SFE5 has inverse scoring scale.

$SF12BP61=100*(((6-SFE5)-1)/4);$

Source variable(s): SFE5

13.4 SF12GH61 (V6 SF-12 General Health Scale)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Note: SFE1 has inverse, renumbered scoring scale.

If SFE1 IN (1,4,5) then $SFE1_ren=6-SFE1;$
If SFE1 IN (2,3) then $SFE1_ren=(6-SFE1)+0.4;$
 $SF12GH61=100*((SFE1_ren-1)/4);$

Source variable(s): SFE1

13.5 SF12VT61 (V6 SF-12 Vitality Scale)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Note: SFE6b has inverse scoring scale.

$$\text{SF12VT61} = 100 * (((6 - \text{SFE6b}) - 1) / 4);$$

Source variable(s): SFE6b

13.6 SF12SF61 (V6 SF-12 Social Functioning Scale)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: $\text{SF12SF61} = 100 * ((\text{SFE7} - 1) / 4);$

Source variable(s): SFE7

13.7 SF12RE61 (V6 SF-12 Role Emotional Scale)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: $\text{SF12RE61} = 100 * (((\text{SFE4a} + \text{SFE4b}) - 2) / 8);$

Source variable(s): SFE4a, SFE4b

13.8 SF12MH61 (V6 SF-12 Mental Health Scale)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: $\text{SF12MH61} = 100 * (((6 - \text{SFE6a}) + \text{SFE6c}) - 2) / 8);$

Source variable(s): SFE6a, SFE6c

13.9 SF12PFZ61 (V6 SF-12 Physical Functioning Scale Z Transformation)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale scores are transformed to Z-scores using US general population means and standard deviations, not age/gender-based.

$$\text{SF12PFZ61} = (\text{SF12PF61} - 81.18122) / 28.10588 ;$$

Source variable(s): SF12PF61

13.10 SF12RPZ61 (V6 SF-12 Role Physical Scale Z Transformation)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale scores are transformed to Z-scores using US general population means and standard deviations, not age/gender-based.

$$\text{SF12RPZ61} = (\text{SF12RP61} - 80.52856) / 27.13526 ;$$

Source variable(s): SF12RP61

13.11 SF12BPZ61 (V6 SF-12 Bodily Pain Scale Z Transformation)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale scores are transformed to Z-scores using US general population means and standard deviations, not age/gender-based.

$$\text{SF12BPZ61} = (\text{SF12BP61} - 81.74015) / 24.53019$$

Source variable(s): SF12BP61

13.12 SF12GHZ61 (V6 SF-12 General Health Scale Z Transformation)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale scores are transformed to Z-scores using US general population means and standard deviations, not age/gender-based.

$$\text{SF12GHZ61} = (\text{SF12GH61} - 72.19795) / 23.19041$$

Source variable(s): SF12GH61

13.13 SF12VTZ61 (V6 SF-12 Vitality Scale Z Transformation)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale scores are transformed to Z-scores using US general population means and standard deviations, not age/gender-based.

$$\text{SF12VTZ61} = (\text{SF12VT61} - 55.59090) / 24.84380$$

Source variable(s): SF12VT61

13.14 SF12SFZ61 (V6 SF-12 Social Functioning Scale Z Transformation)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale scores are transformed to Z-scores using US general population means and standard deviations, not age/gender-based.

$$\text{SF12SFZ61} = (\text{SF12SF61} - 83.73973) / 24.75775 ;$$

Source variable(s): SF12SF61

13.15 SF12REZ61 (V6 SF-12 Role Emotional Scale Z Transformation)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale scores are transformed to Z-scores using US general population means and standard deviations, not age/gender-based.

$$\text{SF12REZ61}=(\text{SF12RE61}-86.41051)/22.35543;$$

Source variable(s): SF12RE61

13.16 SF12MHZ61 (V6 SF-12 Mental Health Scale Z Transformation)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale scores are transformed to Z-scores using US general population means and standard deviations, not age/gender-based.

$$\text{SF12MHZ61}=(\text{SF12MH61}-70.18217)/20.50597;$$

Source variable(s): SF12MH61

13.17 SF12PFT61 (V6 SF-12 Physical Functioning Scale T Score)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale Z-scores are transformed to T-scores.

$$\text{SF12PFT61}=50+(\text{SF12PFZ61}*10);$$

Source variable(s): SF12PFZ61

13.18 SF12RPT61 (V6 SF-12 Role Physical Scale T Score)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale Z-scores are transformed to T-scores.

$$\text{SF12RPT61}=50+(\text{SF12RPZ61}*10);$$

Source variable(s): SF12RPZ61

13.19 SF12BPT61 (V6 SF-12 Bodily Pain Scale T Score)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale scores are transformed to Z-scores using US general population means and standard deviations, not age/gender-based.

$$\text{SF12BPT61} = 50 + (\text{SF12BPZ61} * 10);$$

Source variable(s): SF12BPZ61

13.20 SF12GHT61 (V6 SF-12 General Health Scale T Score)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale Z-scores are transformed to T-scores.

$$\text{SF12GHT61} = 50 + (\text{SF12GHZ61} * 10);$$

Source variable(s): SF12GHZ61

13.21 SF12VTT61 (V6 SF-12 Vitality Scale T Score)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale Z-scores are transformed to T-scores.

$$\text{SF12VTT61} = 50 + (\text{SF12VTZ61} * 10);$$

Source variable(s): SF12VTZ61

13.22 SF12SFT61 (V6 SF-12 Social Functioning Scale T Score)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale Z-scores are transformed to T-scores.

$$\text{SF12SFT61} = 50 + (\text{SF12SFZ61} * 10);$$

Source variable(s): SF12SFZ61

13.23 SF12RET61 (V6 SF-12 Role Emotional Scale T Score)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale Z-scores are transformed to T-scores.

$$\text{SF12RET61} = 50 + (\text{SF12REZ61} * 10);$$

Source variable(s): SF12REZ61

13.24 SF12MHT61 (V6 SF-12 Mental Health Scale T Score)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Scale Z-scores are transformed to T-scores.

$$\text{SF12MHT61} = 50 + (\text{SF12MHZ61} * 10);$$

Source variable(s): SF12MHZ61

13.25 SF12AGGPHYS61 (V6 SF-12 Aggregate Physical Health Score)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Aggregate physical health score created by multiplying scale Z-scores by varimax-rotated factor scoring coefficients and sum the products.

$$\begin{aligned} \text{SF12AGGPHYS61} = & \\ & (\text{SF12PFZ61} * 0.42402) + \\ & (\text{SF12RPZ61} * 0.35119) + \\ & (\text{SF12BPZ61} * 0.31754) + \\ & (\text{SF12GHZ61} * 0.24954) + \\ & (\text{SF12VTZ61} * 0.02877) + \\ & (\text{SF12SFZ61} * -.00753) + \end{aligned}$$

(SF12REZ61* -.19206) +
(SF12MHZ61* -.22069);

Source variable(s): SF12PFZ61, SF12RPZ61, SF12BPZ61, SF12GHZ61, SF12VTZ61,
SF12SFZ61, SF12REZ61, SF12MHZ61

13.26 SF12AGGMENT61 (V6 SF-12 Aggregate Mental Health Score)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Aggregate mental health score created by multiplying scale Z-scores by varimax-rotated factor scoring coefficients and sum the products.

SF12AGGMENT61=
(SF12PFZ61* -.22999) +
(SF12RPZ61* -.12329) +
(SF12BPZ61* -.09731) +
(SF12GHZ61* -.01571) +
(SF12VTZ61* 0.23534) +
(SF12SFZ61* 0.26876) +
(SF12REZ61* 0.43407) +
(SF12MHZ61* 0.48581);

Source variable(s): SF12PFZ61, SF12RPZ61, SF12BPZ61, SF12GHZ61, SF12VTZ61,
SF12SFZ61, SF12REZ61, SF12MHZ61

13.27 SF12AGGPHYST61 (V6 SF-12 Aggregate Physical Health T Score)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Transform Aggregate score to T-score.

SF12AGGPHYST61== 50 + (SF12AGGPHYS61* 10);

Source variable(s): SF12AGGPHYS61

13.28 SF12AGGMENTT61 (V6 SF-12 Aggregate Mental Health T Score)

Format: Continuous variable (f6.2)

Type: Numeric

Algorithm: Transform Aggregate score to T-score.

SF12AGGMENTT61== 50 + (SF12AGGMENT61* 10);

Source variable(s): SF12AGGMENT61