

ARIC Manuscript Proposal # 813S

PC Reviewed: 07/03/01
SC Reviewed: _____

Status: A
Status: _____

Priority: N/A
Priority: _____

SHHS Manuscript Proposal

Full Title:

Susceptibility to sleepiness: factors associated with absence of sleepiness and fatigue in moderate to severe sleep-disordered breathing

Abbreviated Title:

SDB without symptoms

Lead Author:

Vishesh Kapur MD, MPH

Timeline:

Start analysis immediately

Rationale:

Obstructive sleep apnea is defined as the presence of significant SDB accompanied by complaints resulting from disrupted sleep including sleepiness and fatigue. There appear to be individual differences in susceptibility to and/or awareness of sleepiness. It is not uncommon to encounter patients who have significant SDB, but little or no subjective complaints of sleepiness or fatigue. In the Wisconsin sleep cohort only a minority of men and women with RDI>5 had significant sleepiness.¹ Analyses by Gottlieb on the SHHS cohort show that only a minority of individuals with severe SDB have Epworth scores > 10.²

There are studies that have attempted to identify the predictors of objective and subjective sleepiness in sleep clinic patients.^{3,4,5,6,7} These studies indicate that sleep fragmentation, frequency of SDB, female gender are associated with sleepiness while hypoxemia does not appear to be an important factor. Gottlieb's analyses of snoring and sleepiness in the SHHS cohort indicates that RDI is associated with Epworth score and that snoring is an independent predictor of sleepiness at all levels of SDB.^{2,8} In addition, gender, usual sleep duration, sleep restriction on weekdays and symptoms of leg movements were associated with the Epworth score though they did not confound the relationship between the Epworth score and RDI or snoring

Many patients with sleep apnea complain of fatigue in addition to or instead of sleepiness. One study in a sleep laboratory population, patients more frequently complained of tiredness than sleepiness.⁹ There are no community-based studies that have determined the prevalence of symptoms of fatigue relative to sleepiness in moderate to severe SDB. In addition, there are no community-based studies that have identified factors that are associated with the absence of fatigue and sleepiness in individuals with moderate to severe SDB.

The identification of factors associated with the *absence* of sleepiness and fatigue may important to identify individuals with significant SDB who do not have these complaints but still have increased

risk for the potential medical complications of SDB. In addition, the identification of factors associated with sleepiness and fatigue in individuals with significant SDB may provide insights into individual susceptibility to sleepiness and the causal relationship between SDB and symptoms of non-restorative sleep. The SHHS has collected data that can address these issues.

Hypotheses:

1. A majority of subjects with moderate to severe SDB do not have complaints of sleepiness or fatigue. Fatigue is a more common complaint than sleepiness in this group.
2. The absence of fatigue and sleepiness in this population is associated with:
 - a. Less disrupted sleep as assessed by EEG
 - b. Longer usual sleep duration
 - c. No evidence of weekday sleep restriction
 - d. Female gender
 - e. Lower educational level
 - f. Lack of use of sedating medications
 - g. Absence of self-report of awakenings due to legs
 - h. Less frequent snoring
 - i. Less comorbid illness

Data:

PSG: RDI, Arousal Index, Awakening Index, sleep efficiency, sleep stage %, lowest oxygen saturation, % time spent below 90% saturation

Sleep habits questionnaire: Questions on fatigue, sleepiness, sleep hours, snoring leg jerks

SF-36: Question on frequency of tiredness

Medications: Use of MAOI, TCA, benzodiazepine, antipsychotics. The chronic disease score calculated from medication list (see manuscript #27)

Demographic: Age, gender, BMI, race, education level

General Health: smoking caffeine, alcohol, self-reported cardiovascular and pulmonary disease

Type of Study:

Case-control study

Type of Publication:

Journal article

Analysis Responsibility:

Author

Introduction:

These analyses will determine the proportion of subjects with moderate to severe SDB ($RDI \geq 15$) that have fatigue, sleepiness, both or neither and which factors are associated with the absence of symptoms.

Brief Analysis Plan:

1. Determine the prevalence of sleepiness, fatigue, both symptoms and no symptoms in subject with $RDI \geq 15$ using questions from sleep habits (#7E) and SF-36 (9i) on frequency of these symptoms. Compare characteristics of subjects with fatigue only, sleepiness only, both symptoms and no symptoms
2. Case control study:

Cases: $RDI \geq 15$ with rare or less frequent sleep and “a little of the time” or less frequently tired

Controls: $RDI \geq 15$ with at least often sleep or “a good bit of the time” tired

Independent variables: listed under data

Primary statistical method: logistic regression

Compare cases to controls and estimate odds ratios for factors associated sleepiness or fatigue.

Identify and estimate odds ratios for independent risk factors using multivariate models.

Summary: The results of these analyses will add to our understanding of how individuals with moderate SDB perceive the effects of sleep disruption and what factors are associated with an awareness of the effects.

1. Young T et al. The occurrence of sleep-disordered breathing among middle-aged adults. *New Eng J Med* 1993; 328(17): 1230-5.
2. Gottlieb DJ et al. Relation of sleepiness to respiratory disturbance index: the Sleep Heart Health Study. *Am J Respir Crit Care Med* 1999 Feb; 159(2): 502-7.
3. Guilleminault C et al. Determinants of daytime sleepiness in obstructive sleep apnea. *Chest* 1988 Jul; 94(1):32-7.
4. Roehrs T et al. Predictors of objective level of daytime sleepiness in patients with sleep-related breathing disorders. *Chest* 1989 Jun; 95(6): 1202-6.
5. Furuta H et al. Epworth Sleepiness Scale and sleep studies in patients with obstructive sleep apnea syndrome. *Psychiatry Clin Neurosci* 1999 Apr; 53(2): 301-2.
6. Cot HG et al. Hypoxemia vs. sleep fragmentation as cause of excessive daytime sleepiness in obstructive sleep apnea. *Chest* 1991 Dec; 100(6): 1542-8.
7. Johns MW. Daytime sleepiness, snoring and obstructive sleep apnea. The Epworth Sleepiness Scale. *Chest* 1993 Jan; 103(1): 30-6.
8. Gottlieb DJ et al. Does snoring predict sleepiness independently of apnea and hypopneas frequency? *Am J Respir Crit Care Med*. 162: 1512-1517.
9. Chervin RD. Sleepiness, fatigue, tiredness and lack of energy in obstructive sleep apnea. *Chest* 2000 Aug; 118(2): 372-9.